

---

**MONTANA DEPARTMENT OF TRANSPORTATION  
WETLAND MITIGATION MONITORING REPORT: YEAR 2006**

---

*Beaverhead Gateway  
Dillon, Montana*



Prepared for:

**MONTANA DEPARTMENT OF TRANSPORTATION**  
2701 Prospect Ave  
Helena, MT 59620-1001

Prepared by:

**POST, BUCKLEY, SCHUH, & JERNIGAN**  
P.O. Box 239  
Helena, MT 59624

December 2006

Project No: B43054.00 - 0202



# **MONTANA DEPARTMENT OF TRANSPORTATION**

## **WETLAND MITIGATION MONITORING REPORT:**

**YEAR 2006**

*Beaverhead Gateway  
Dillon, Montana*

Prepared for:

**MONTANA DEPARTMENT OF TRANSPORTATION**  
2701 Prospect Ave  
Helena, MT 59620-1001

Prepared by:

**POST, BUCKLEY, SCHUH, & JERNIGAN**  
P.O. Box 239  
Helena, MT 59624

December 2006

Project No: B43054.00 - 0202

***“MDT attempts to provide accommodations for any known disability that may interfere with a person participating in any service, program, or activity of the Department of Transportation. Alternative accessible formats of this information will be provided upon request. For further information, call 406-444-7228 or TTY (800-335-7592) or by calling Montana Relay at 711.”***



## TABLE OF CONTENTS

<b>1.0 INTRODUCTION.....</b>	<b>1</b>
<b>2.0 METHODS.....</b>	<b>3</b>
2.1 Monitoring Dates and Activities.....	3
2.2 Hydrology .....	3
2.3 Vegetation.....	3
2.4 Soils.....	4
2.5 Wetland Delineation .....	4
2.6 Mammals, Reptiles and Amphibians .....	4
2.7 Birds.....	4
2.8 Macroinvertebrates .....	5
2.9 Functional Assessment.....	5
2.10 Photographs.....	5
2.11 GPS Data.....	5
2.12 Maintenance Needs.....	5
<b>3.0 RESULTS .....</b>	<b>5</b>
3.1 Hydrology .....	5
3.2 Vegetation.....	6
3.3 Soils.....	12
3.4 Wetland Delineation .....	13
3.5 Wildlife .....	13
3.6 Macroinvertebrates .....	15
3.7 Functional Assessment.....	16
3.8 Photographs.....	17
3.9 Maintenance Needs/Recommendations .....	17
3.10 Current Credit Summary.....	18
<b>4.0 REFERENCES.....</b>	<b>19</b>

## **TABLES**

Table 1	<i>2001-2006 vegetation species list for the Beaverhead Gateway Mitigation Site.</i>
Table 2	<i>Transect 1 data summary.</i>
Table 3	<i>Transect 2 data summary.</i>
Table 4	<i>2006 Wetland conditions within the Beaverhead Gateway Wetland Mitigation Site.</i>
Table 5	<i>Fish and wildlife species observed at the Beaverhead Gateway Wetland Mitigation Site from 2001 to 2006.</i>
Table 6	<i>Summary of 2006 wetland function/value ratings and functional points for the Beaverhead Gateway Wetland Mitigation Site.</i>

## **FIGURES**

Figure 1	<i>Project Site Location Map</i>
Figure 2	<i>2006 Monitoring Activity Locations</i>
Figure 3	<i>2006 Mapped Site Features</i>

## **CHARTS**

Chart 1	<i>Transect maps showing vegetation types for Transect 1 from start (0 feet) to end (1650 feet) for each year monitored.</i>
Chart 2	<i>Length of vegetation community types along Transect 1 for each year monitored.</i>
Chart 3	<i>Transect maps showing vegetation types for Transect 2 from start (0 feet) to end (280 feet) for each year monitored.</i>
Chart 4	<i>Length of vegetation community types within Transect 2 for each year monitored.</i>
Chart 5	<i>Bioassessment scores for Beaverhead Gateway Wetland Mitigation Site from 2001 to 2006.</i>

## APPENDICES

Appendix A *Figures 2 & 3*

Appendix B *2006 Wetland Mitigation Site Monitoring Form*

*2006 Wetland Delineation Forms*

*2006 Bird Survey Forms*

*2006 COE Wetland Delineation Forms*

*2006 MDT Functional Assessment Forms*

Appendix C *2006 Representative Photographs*

*2001-2006 Aerial Photograph Comparison*

Appendix D *Original Site Plan*

*MDT Bird Observations*

*Letters Addressing Site Management*

Appendix E *Bird Survey Protocol*

*GPS Protocol*

Appendix F *2006 Macroinvertebrate Sampling Protocol and Data Analysis*

## 1.0 INTRODUCTION

This report represents the sixth and final year of monitoring at the Beaverhead Gateway Ranch wetland mitigation site by Post, Buckley, Schuh, and Jernigan (PBS&J). The Beaverhead Gateway Ranch wetland mitigation site was developed to mitigate wetland impacts associated with Montana Department of Transportation (MDT) roadway projects in Watershed 6 located in the MDT Butte District. Some of these projects are completed and some have yet to be constructed. The mitigation site is located within a 196-acre conservation easement 13 miles northeast of Dillon and 14 miles southwest of Twin Bridges on Highway 41 (**Figure 1**). Elevations range from approximately 4825 to 4830 feet. The western portion of the site is in Beaverhead County and the eastern portion is in Madison County. MDT personnel monitored the site in 1998, 1999 and 2000.

The approximate site boundary is illustrated on **Figure 2** in **Appendix A**, and the original site plans are included in **Appendix D**. The project is located adjacent to the Beaverhead River and Highway 41. Upwelling groundwater and springs with surface retention behind a constructed dike provides wetland hydrology. Precipitation and surface runoff provide minor contributions to wetland hydrology at this site. The site is in private ownership and occurs within a conservation easement. The wetland easement area is not fenced exclusively; however, portions of the easement are fenced for cattle management (through the Montana State University livestock extension program) and the larger property containing the easement is fenced.

Construction was completed in 1997 with the goal of creating at least 52 acres of wetland. The site includes a dike constructed to retain storm water and groundwater collected in two prior-existing drainage ditch systems. A control structure was completed in the northwest portion of the impoundment located where the two former drainage ditches converged. This control structure can be used to adjust impoundment water levels. The impoundment was designed to inundate approximately 26 acres with water depths of 0 to 3 feet.

The site was designed to mitigate for specific wetland functions impacted by MDT roadway projects, including: storm water retention, roadway runoff filtration, sediment and nutrient retention, water quality, groundwater recharge, waterfowl and wildlife habitats and riparian restoration. In addition to creating 52 acres of new wetland, a primary goal is to use an ephemeral creek channel entering the southeastern quadrant of the site to capture storm water flows from nearby farmland and allow silts/suspended sediments to settle out within the wetland.

A pre-project construction wetland delineation documented 5.2 acres of wetlands at the site (Hackley 1997). The monitoring area is illustrated in **Figure 2** in **Appendix A**.



## 2.0 METHODS

### 2.1 Monitoring Dates and Activities

The site was visited on June 1 (spring season), August 9 (mid-season) and September 17, 2006 (fall season). The spring and fall visits were conducted to sample seasonal bird and other wildlife uses. Spring season monitoring is likely to detect migrant and early nesting activities for a variety of avian species (Carlson pers. comm.), as well as maximize the potential for amphibian detection. In Montana, most amphibian larval stages are present by early June (Werner pers. comm.).

The mid-season visit was conducted in July to document vegetation, soil, and hydrologic conditions used to map jurisdictional wetlands. All information contained on the Wetland Mitigation Site Monitoring Form (**Appendix B**) was collected at this time. Activities and information conducted/collected included: wetland delineation; wetland/open water aquatic habitat boundary mapping; vegetation community mapping; vegetation transect; soils data; hydrology data; bird and general wildlife use; photograph points; macroinvertebrate sampling; GPS data points; functional assessment; and (non-engineering) examination of dike structures.

### 2.2 Hydrology

Wetland hydrology indicators were recorded during the mid-season visit using procedures outlined in the COE 1987 Wetland Delineation Manual (Environmental Laboratory 1987). Hydrology data were recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**).

Additional hydrologic data were recorded on the mitigation site monitoring form (**Appendix B**). Although two deep remnant wells remain on the property in the wetland vicinity (which were not sampled), no groundwater monitoring wells were installed at the site; consequently, no groundwater monitoring was conducted. If present within 18 inches of the ground surface (soil pit depth for purposes of delineation), groundwater depths were documented on the routine wetland delineation data form at each data point.

### 2.3 Vegetation

General dominant species-based vegetation community types (e.g., *Alopecurus/Juncus*) were delineated on an aerial photograph during the mid-season visit. Standardized community mapping was not employed as many of these systems are geared towards climax vegetation and do not reflect yearly changes. Estimated percent cover of the dominant species in each community type was listed on the Wetland Mitigation Site Monitoring Form (**Appendix B**).

Two 10-foot wide belt transects established in 2001 were sampled during the mid-season monitoring event to represent the range of current vegetation conditions. Percent cover was estimated for each vegetative species within each successive vegetative community encountered within the “belt” using the following values: T (few plants); P (1-5%), 1 (5-15%); 2 (15-25%); 3 (25-35%); 4 (35-45%); 5 (45-55%) and so on to 9 (85-95%). The transect locations are illustrated on **Figure 2** in **Appendix A**. These transects are used to evaluate changes over time,

especially the establishment and increase of hydrophytic vegetation. The transect data were recorded on the mitigation site monitoring form, and photographs were taken from both ends of each transect looking along the transect path.

A comprehensive plant species list for the site was updated as new species were encountered. Ultimately, observations from past years will be compared with new data to document vegetation changes over time. Woody species were not planted at this mitigation site.

## 2.4 Soils

Soils were evaluated during the mid-season site visit using the hydric soils determination procedures outlined in the COE 1987 Wetland Delineation Manual. Soil data were recorded for each wetland determination point on the COE Routine Wetland Delineation Data Forms (**Appendix B**). The most current terminology used by NRCS was used to describe hydric soils (USDA 1998).

## 2.5 Wetland Delineation

Wetland delineation was conducted during the mid-season visit according to the 1987 COE Wetland Delineation Manual. Wetland and upland areas within the monitoring area were investigated for the presence of wetland hydrology, hydrophytic vegetation and hydric soils. The information was recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**). The wetland/upland boundary was originally delineated on the aerial photo and recorded with a resource grade GPS unit in 2001 using procedures outlined in **Appendix E**. Modifications to these boundaries in 2006 were accomplished by hand-mapping onto the 2006 aerial photograph. The wetland/upland boundary in combination with the wetland/open water boundary was used to calculate the final wetland acreage. A pre-construction wetland delineation documented 5.2 acres of wetlands at the site (Hackley 1997).

## 2.6 Mammals, Reptiles, and Amphibians

Mammal, reptile, and amphibian species observations and other positive indicators of use, such as vocalizations, were recorded on the wetland monitoring and bird forms during the 2006 monitoring events. Indirect use indicators, including tracks, scat, burrows, eggshells, skins, bones, etc. were also recorded. Observations were recorded as the observer traversed the site while conducting other required activities. Direct sampling methods, such as snap traps, live traps, and pitfall traps, were not used.

## 2.7 Birds

Bird observations were recorded during the spring, mid-season, and fall visits. No formal census plots, spot mapping, point counts, or strip transects were conducted. The spring and fall birding visit was conducted in accordance with the Bird Survey Protocols (**Appendix E**). During the mid-season visit, the bird survey was conducted incidental to other monitoring activities. Bird species observations were categorized by species, activity code, and general habitat association on the Bird Survey Field Data Sheet (**Appendix B**).

## 2.8 Macroinvertebrates

Macroinvertebrate samples were collected during the mid-season site visit at four separate locations (**Figure 2** in **Appendix A**). In previous years two additional samples were collected for a total of six. However, in 2004 to 2006 water was absent at macroinvertebrate sampling site 2 and 4, and no collections were made. Collections occurred using the Macroinvertebrate Sampling Protocol (**Appendix F**). Samples were preserved as outlined in the sampling procedure and sent to Rhithron Associates, Inc. in Missoula, Montana for analysis (**Appendix F**).

## 2.9 Functional Assessment

A functional assessment form was completed using the 1999 MDT Montana Wetland Assessment Method (Berglund 1999) (**Appendix B**). Field data necessary for this assessment were collected during the mid-season visit. No pre-project functional assessment was conducted at this site.

## 2.10 Photographs

Photographs were taken illustrating current land uses surrounding the site, the upland buffer, the monitored area and the vegetation transects (**Appendix C**). Each photograph point location was recorded with a resource grade GPS in 2001. The location of photo points is shown on **Figure 2** in **Appendix A**. All photographs were taken using a digital camera.

## 2.11 GPS Data

During the 2001 monitoring season, point data were collected with a resource grade GPS unit at the vegetation transect beginning and ending locations and at all photograph locations. Wetland boundaries were also recorded with a resource grade GPS unit in 2001, but were modified via hand-mapping onto an aerial photograph in subsequent years. Procedures used for GPS mapping and aerial photograph referencing are included in **Appendix E**.

## 2.12 Maintenance Needs

Observations were made of existing structures and of erosion/sediment problems to identify maintenance needs. This did not constitute an engineering-level structural inspection, but rather a cursory examination. Current or future potential problems were documented on the monitoring form.

## 3.0 RESULTS

### 3.1 Hydrology

The main source of hydrology seems to be upwelling groundwater and “springs” evident along the constructed channels (ditch/berms) leading south and west from the main open water area (**Figure 3** in **Appendix A**). Water was observed upwelling from the bottom of these channels.

These waters are retained behind a constructed dike. Another source of hydrology comes from the southeast corner of the site from irrigation return flow. Precipitation and surface runoff provide minor contributions to wetland hydrology at this site except during rare and extreme events.

Open water occurred across approximately 6.5 acres or 5% of the 118-acre wetland area (**Figure 3** in **Appendix A**) during the mid-season visit. Water depth at the open water/rooted vegetation boundary was approximately 1.5 feet. Inundation was observed during the mid-season visit across approximately another 5% of the wetland area which is similar to the last three years (2003-2005) but significantly less than in the two preceding years (2001 and 2002). Inundation was present throughout most of Community Type 2 (**Figure 3** in **Appendix A**), and portions of Type 8. Casual observations during the early season visit indicated complete inundation of Types 2 and 8 and more extensive inundation throughout Type 6.

Annual precipitation totals for 2004, 2005 and 2006 were 10.82, 12.36 and 9.14 inches (to date), respectively, compared with the long term annual mean of 9.77 inches. Precipitation from January through July in 2006 was 6.78 inches, which exceeded the long-term January-July mean of 6.55 inches. Despite this, water levels in 2004, 2005 and 2006 at the site were lower than observed during past monitoring in 2001 and 2002, when annual precipitation totaled 6.82 and 9.17 inches, respectively. The reason for lower water levels over the past three years is not clear, but may be related to delayed effects of reduced precipitation from 1999 – 2003 during which annual precipitation levels were generally well below the long-term mean, allowing for greater influence of evapotranspiration, percolation, and leakage. Also, in 2004, the landowner dropped the water level by removing some stop logs in order to reduce wave action along the main dike and facilitate fabric and gravel installation. While this activity may have temporarily affected 2004 water levels, it is unlikely to have substantively affected 2006 levels. Another reason for lower observed water levels may relate to possible changed irrigation practices up-gradient. As requested by MDT, letters discussing water level management related to dike erosion issues are included in **Appendix D**.

None of the six wetland sites documented on the Routine Wetland Determination forms (**Appendix B**) had groundwater within 18 inches of the surface on August 8, 2006. Casual observations at other locations on this date revealed groundwater within 18 inches of the surface in small areas of Community Types 2 and 6 (**Figure 3** in **Appendix A**). These groundwater depths seem low compared with the soil and vegetation indicators present and are similar to depths observed in 2005. Continued low groundwater depths could result in a decline in wetland vegetation. It is important to note that drought conditions have dominated for many years in recent time. Hydrologic conditions must be considered within this climatic context.

### 3.2 Vegetation

Ninety-seven plant species were identified at the site and are listed in **Table 1**. No new species were identified in 2006. The majority of these species were herbaceous. Few woody species were found within the monitoring area. One plant species of concern, Lemmon's Alkali Grass (*Puccinellia lemmonii*), was identified in past years and is ranked S1 by the Montana Natural Heritage Program. However, Lemmon's Alkali Grass was not observed in 2006.

**Table 1: 2001-2006 vegetation species list for the Beaverhead Gateway Mitigation Site.**

Scientific Name	Common Name	Region 9 (Northwest) Wetland Indicator
<i>Agropyron cristatum</i>	crested wheatgrass	--
<i>Agropyron repens</i>	quack grass	FACU
<i>Agropyron smithii</i>	western wheatgrass	FACU
<i>Agropyron trachycaulum</i>	slender wheatgrass	FAC
<i>Agrostis stolonifera</i>	redtop	FAC+
<i>Alopecurus pratensis</i>	meadow foxtail	FACW
<i>Artemisia frigida</i>	fringed sagewort	--
<i>Artemisia</i> spp.	sagebrush	--
<i>Aster falcatus</i>	leafy-bracted aster	FACU-
<i>Aster hesperius</i>	Siskiyou aster	OBL
<i>Astragalus</i> spp.	milkvetch	--
<i>Bromus inermis</i>	smooth brome	--
<i>Bromus japonicus</i>	Japanese brome	FACU
<i>Bromus tectorum</i>	cheatgrass	--
<i>Calamagrostis neglecta</i>	slim reedgrass	FACW
<i>Cardaria draba</i>	white top	--
<i>Carduus nutans</i> *	musk thistle	--
<i>Carex capillaries</i>	hair-like sedge	FACW
<i>Carex limnophila</i>	pond sedge	FACW
<i>Carex nebrascensis</i>	Nebraska sedge	OBL
<i>Carex praegracilis</i>	clustered field sedge	FACW
<i>Carex torreyi</i> *	Torrey's sedge	FAC
<i>Centaurea maculosa</i> *	spotted knapweed	--
<i>Chenopodium album</i>	white goosefoot	FAC
<i>Chenopodium rubrum</i>	coastal-blite pigweed	FACW+
<i>Chrysothamnus nauseosus</i>	rubber rabbitbrush	--
<i>Cirsium arvense</i>	Canada thistle	FACU+
<i>Cirsium undulatum</i>	wavy-leaf thistle	FACU+
<i>Cleome serrulata</i>	Rocky Mountain bee plant	FACU
<i>Cornus stolonifera</i> *	red-osier dogwood	FACW
<i>Cynoglossum officinalis</i>	hound's tongue	FACU
<i>Dactylis glomerata</i>	orchard grass	FACU
<i>Descurainia sophia</i>	tansy mustard	--
<i>Distichlis spicata</i>	saltgrass	FAC+
<i>Elaeagnus angustifolia</i> *	Russian olive	FAC
<i>Eleocharis acicularis</i> *	least spike rush	OBL
<i>Eleocharis pauciflora</i>	few-flowered spike rush	OBL
<i>Elymus cinereus</i>	big basin wild rye	FACU
<i>Epilobium palustris</i>	swamp willow-herb	OBL
<i>Equisetum laevigatum</i>	smooth scouring-rush	FACW
<i>Festuca idahoensis</i>	Idaho fescue	FACU
<i>Festuca pratensis</i>	meadow fescue	FACU+
<i>Gentianella amarelle</i>	northern gentian	FACW-
<i>Glaux maritime</i>	sea-milkwort	FACW+
<i>Grindelia squarrosa</i>	curly-cup gumweed	FACU
<i>Habenaria dilatata</i>	bog orchid	--
<i>Haplopappus carthamoides</i>	Columbia goldenweed	--
<i>Helianthus nuttalli</i>	Nuttall's sunflower	FACW-
<i>Helenium autumnale</i> *	sneezeweed	FACW
<i>Hippuris vulgaris</i>	common mare's-tail	OBL
<i>Hordeum jubatum</i>	foxtail barley	FAC+
<i>Iris missouriensis</i>	Rocky Mountain iris	OBL
<i>Iva axillaries</i>	small-flower sumpweed	FAC
<i>Juncus balticus</i>	Baltic rush	FACW+
<i>Juncus bufonius</i>	toad rush	FACW+

**Table 1 (continued): 2001-2006 vegetation species list for the Beaverhead Gateway Mitigation Site.**

Scientific Name	Common Name	Region 9 (Northwest) Wetland Indicator
<i>Juncus ensifolius</i>	three-stamen rush	FACW
<i>Kochia scoparia</i>	summer-cypress	FAC
<i>Lactuca serriola</i>	prickly lettuce	FAC-
<i>Lepidium perfoliatum</i>	clasping pepper-grass	FACU+
<i>Lycopus asper</i>	rough bugleweed	OBL
<i>Medicago lupulina</i>	black medic	FAC
<i>Medicago sativa</i>	alfalfa	--
<i>Melilotus alba</i>	white sweetclover	FACU
<i>Melilotus officinalis</i>	yellow sweetclover	FACU
<i>Mentha arvensis</i> *	mint	FAC
<i>Mimulus</i> spp.*	monkey flower	OBL
<i>Muhlenbergia asperifolia</i>	alkali muhly	FACW
<i>Myosotis discolor</i> *	forget-me-not	FACW
<i>Phalaris arundinacea</i>	canary reed grass	FACW
<i>Phleum pratense</i> *	timothy	FACU
<i>Plantago eriopoda</i>	saline plantain	FACW
<i>Phlox longifolia</i>	long-leaf phlox	--
<i>Phragmites australis</i> *	common reed	FACW+
<i>Poa pratensis</i>	Kentucky bluegrass	FACU+
<i>Poa sandbergii</i>	Sandberg's bluegrass	--
<i>Polygonum amphibium</i> *	water smartweed	OBL
<i>Polygonum aviculare</i>	prostrate knotweed	FACW+
<i>Populus trichocarpa</i> *	cottonwood	FAC
<i>Potamogeton</i> spp.*	pondweed	OBL
<i>Potentilla anserine</i>	silverweed	OBL
<i>Potentilla fruticosa</i> *	shrubby cinquefoil	FAC-
<i>Puccinellia lemmonii</i>	Lemmons alkali grass	FAC
<i>Ranunculus populago</i>	poplar buttercup	FACW
<i>Rorippa</i> spp.*	watercress	OBL
<i>Rumex crispus</i> *	curly dock	FACW
<i>Salicornia</i> spp.*	saltwort	--
<i>Salix bebbiana</i> *	Bebb willow	FACW
<i>Salix exigua</i>	sandbar willow	OBL
<i>Salsola kali</i>	Russian thistle	FACU
<i>Sarcobatus vermiculatus</i>	greasewood	FACU+
<i>Scirpus acutus</i> *	hard stem bulrush	OBL
<i>Scirpus americanus</i>	American bulrush	OBL
<i>Scirpus maritimus</i> *	salt marsh bulrush	OBL
<i>Scirpus pungens</i>	three-square bulrush	OBL
<i>Scirpus validus</i>	soft-stem bulrush	OBL
<i>Shepherdia</i> spp.*	buffaloberry	--
<i>Sisyrinchium angustifolium</i>	western blue eyed grass	FACW-
<i>Sonchus arvensis</i>	field sowthistle	FAC-
<i>Spartina gracilis</i>	alkali cordgrass	FACW
<i>Sporobolus cryptandrus</i>	sand dropseed	FACU
<i>Stipa comata</i>	needle & thread grass	--
<i>Suaeda intermedia</i>	alkali seepweed	FAC
<i>Tragopogon dubius</i>	yellow salsify	--
<i>Triglochin maritime</i>	seaside arrowgrass	OBL
<i>Typha latifolia</i>	cattail	OBL
<i>Urtica dioica</i>	stinging nettle	FAC+
<i>Zigadenus venenosus</i>	meadow death camas	FAC

\* - Plant species observed by Montana Department of Transportation.

Four Wetland Community types (Type 2: *Scirpus*, Type 5: *Alopecurus/Juncus*, Type 6: *Alopecurus/Scirpus* and Type 8: *Potamogeton/Polygonum*) and three Upland Community Types (Type 3: *Hordeum/Kochia*, Type 4: *Muhlenbergia/Agropyron* and Type 7: *Sarcobatus/Elymus*) were identified and mapped at the mitigation area (**Figure 3** in **Appendix A**). Plant species observed within each of these communities are listed on the attached data form (**Appendix B**).

Type 8 is the wettest community type and occurred as an aquatic bed community in the shallower water areas (**Figure 3**). It was dominated by pondweed (*Potamogeton spp.*) and smartweed (*Polygonum spp.*). Type 2 is the next wettest and occurred mainly as a fringe around the border of shallow water areas dominated by bulrush (*Scirpus spp.*). Type 6 is the next wettest wetland vegetation type and occurred throughout the monitoring area on sites slightly higher than Type 2. The vegetation in Type 6 was highly variable from spot to spot due to small changes in soil properties, topography, and past disturbance. Vegetation in Type 6 was also highly variable since it was in transition from upland to wetland. Across much of this type, the vegetation was dominated by meadow foxtail (*Alopecurus pratensis*) and bulrush. However, small areas were dominated by other species.


Adjacent upland vegetation community types were mainly dominated by rangeland species with cropland along the southern border. Type 3 was located along dikes, spoil pile and or other highly disturbed soil materials and was dominated by weedy species such as foxtail barley (*Hordeum jubatum*), summer-cypress (*Kochia scoparia*) and Canada thistle (*Cirsium arvense*). Type 4 was mostly dominated by alkali muhly (*Muhlenbergia asperifolia*), slender wheatgrass (*Agropyron trachycaulum*) and western wheatgrass (*Agropyron smithii*). Type 7 was dominated by greasewood (*Sarcobatus vermiculatus*), basin wild rye (*Elymus cinereus*) and western wheatgrass.

Vegetation transect results are detailed in the attached data form (**Appendix B**), and are summarized in **Tables 2** and **3** and **Charts 1** to **4**. Vegetation transects results showed no change between each monitoring year.


**Table 2: Transect 1 data summary.**

Monitoring Year	2001	2002	2003	2004	2005	2006
Transect Length (feet)	1650	1650	1650	1650	1650	1650
# Vegetation Community Transitions along Transect	5	5	5	5	5	5
# Vegetation Communities along Transect	4	4	4	4	4	4
# Hydrophytic Vegetation Communities along Transect	3	3	3	3	3	3
Total Vegetative Species	22	22	22	22	22	22
Total Hydrophytic Species	12	14	14	14	14	14
Total Upland Species	10	8	8	8	8	8
Estimated % Total Vegetative Cover	95	95	95	95	95	95
% Transect Length Comprised of Hydrophytic Vegetation Communities	98	98	98	98	98	98
% Transect Length Comprised of Upland Vegetation Communities	2	2	2	2	2	2
% Transect Length Comprised of Unvegetated Open Water	0	0	0	0	0	0
% Transect Length Comprised of Bare Substrate	0	0	0	0	0	0

**Chart 1: Transect maps showing vegetation types for Transect 1 from start (0 feet) to end (1650 feet) for each year monitored.**




**Chart 2: Length of vegetation community types within Transect 1 for each year monitored.**




**Table 3: Transect 2 data summary.**

Monitoring Year	2001	2002	2003	2004	2005	2006
Transect Length (feet)	280	280	280	280	280	280
# Vegetation Community Transitions along Transect	4	4	4	4	4	4
# Vegetation Communities along Transect	4	4	4	4	4	4
# Hydrophytic Vegetation Communities along Transect	2	2	2	2	2	2
Total Vegetative Species	18	21	21	21	21	22
Total Hydrophytic Species	11	10	10	10	10	10
Total Upland Species	7	11	11	11	11	12
Estimated % Total Vegetative Cover	80	80	80	80	80	89
% Transect Length Comprised of Hydrophytic Vegetation Communities	43	43	43	43	43	43
% Transect Length Comprised of Upland Vegetation Communities	57	57	57	57	57	57
% Transect Length Comprised of Unvegetated Open Water	0	0	0	0	0	0
% Transect Length Comprised of Bare Substrate	0	0	0	0	0	0

**Chart 3: Transect maps showing vegetation types for Transect 2 from start (0 feet) to end (280 feet) for each year monitored.**



**Chart 4: Length of vegetation community types within Transect 2 for each year monitored.**



Noxious weeds at the site included spotted knapweed (*Centaurea maculosa*) and Canada thistle. Other weedy species included summer-cypress, hound's-tongue (*Cynoglossum officinalis*), curly-cup gumweed (*Grindelia squarrosa*), lambsquarters (*Chenopodium album*), whitetop (*Cardaria draba*) and quackgrass (*Agropyron repens*). No common reed (*Phragmites australis*) was observed at the site although it was present nearby along Highway 41. This is an extremely aggressive invader of wetlands and a serious concern at this site. Weed control and revegetation is needed at this site to prevent further spread and protect soil from wind and water erosion. Additional effort should be made to determine if common reed or other important weeds are present.

### 3.3 Soils

The western two-thirds of the site are within Beaverhead County where soil survey information is not currently available. The eastern one-third of the site was mapped as part of the Madison County Soil Survey (USDA 1989). The soil in the eastern one-third of the site is mapped as Neen silty clay loam with randomly distributed soils that have a layer of organic material 4 to 20 inches thick at the surface (USDA 1989). Neen soils are not listed on the Montana NRCS Hydric Soil list. Soil characteristics at each wetland determination point were compared with those of the Neen soil. The soils observed across most of the site did not generally match the Neen soil. The main portion of the site mapped during the Madison County soil survey is currently under water.

Wetland soils were similar to those observed in 2001-2005. Wetland soils observed during monitoring and documented on the Routine Wetland Determination form were mostly loams, silt loams or silty clay loams with very low chromas (0 or 1) within 2 inches of the surface. Mottles (redoximorphic features) were present in most profiles observed. Only one of four soil profiles described on the Routine Wetland Determination forms was saturated within 18 inches of the surface reflecting the time of year and the recent history of drought discussed above. Small areas were observed with thin organic surface layers and with mucky mineral surface layers.

### 3.4 Wetland Delineation

Wetland boundaries were similar in 2006 to those mapped in past years. These wetland boundaries continue to be located at distinct topographic and soil breaks that are not likely to change over time without a significant change in groundwater elevation or climate. Delineated wetland boundaries are illustrated on **Figure 3** in **Appendix A**. Completed wetland delineation forms are included in **Appendix B**. Soils, vegetation, and hydrology are discussed in preceding sections.

Wetland conditions identified from monitoring from 2001 through 2006 are described in **Table 4**. Approximately 111.7 wetland acres and 6.5 open water acres occur within the 2006 monitoring area (**Figure 3** in **Appendix A**). The pre-construction wetland delineation reported 5.2 wetland and no open water acres. The net increase in wetland acres is  $111.7 - 5.2 = 106.5$  acres plus 6.5 acres of open water. Additional area may form with time and more normal precipitation around the low gradient portions of the current wetland area.

**Table 4: 2006 Wetland conditions within the Beaverhead Gate Wetland Mitigation Site.**

Wetland Condition	Monitoring Area	Above Dike	Below Dike
Gross Wetland Area	118.2	97.9	20.3
Open Water Area	6.5	6.5	0.0
<b>Net Wetland Area</b>	<b>111.7</b>	<b>91.4</b>	<b>20.3</b>

### 3.5 Wildlife

Wildlife species, or evidence of wildlife, observed on the site during 2001-2006 monitoring efforts are listed in **Table 5**. The site receives substantial use by American white pelicans, trumpeter swans, black terns, sandhill cranes, and other species. Sandhill cranes are known breeders on the site (Urban pers. comm.). American white pelicans, trumpeter swans, and black terns are all considered species of concern by the MNHP relative to breeding locations. Of these three species, black terns are likely breeders on the site.

**Table 5: Fish and wildlife species observed at the Beaverhead Gateway Mitigation Site from 2001 to 2006<sup>1</sup>.**

<b>FISH</b>	
None	
<b>AMPHIBIAN</b>	
None	
<b>REPTILES</b>	
Garter Snake ( <i>Thamnophis</i> spp.) Painted Turtle	
<b>BIRDS</b>	
<p><b>American White Pelican</b> (<i>Pelecanus erythrorhynchos</i>)</p> <p>American Crow (<i>Corvus brachyrhynchos</i>)</p> <p><b>American Coot</b> (<i>Fulica americana</i>)</p> <p>American Dipper (<i>Cinclus</i>)</p> <p><b>Bald Eagle</b> (<i>Haliaeetus leucocephalus</i>)</p> <p>Bank Swallow (<i>Riparia riparia</i>)</p> <p><b>Black-billed Magpie</b> (<i>Pica pica</i>)</p> <p>Black-necked Stilt (<i>Himantopus mexicanus</i>)</p> <p>Black Tern (<i>Chlidonias niger</i>)</p> <p><b>Blue-winged Teal</b> (<i>Anas discors</i>)</p> <p>Bullock's Oriole (<i>Icterus bullockii</i>)</p> <p><b>Canada Goose</b> (<i>Branta canadensis</i>)</p> <p><b>Cinnamon Teal</b> (<i>Anas cyanoptera</i>)</p> <p><b>Cliff Swallow</b> (<i>Petrochelidon pyrrhonota</i>)</p> <p><b>Common Goldeneye</b> (<i>Bucephala clangula</i>)</p> <p>Common Snipe (<i>Gallinago gallinago</i>)</p> <p>Common Yellowthroat (<i>Geothlypis trichas</i>)</p> <p>Cowbird (<i>Molothrus ater</i>)</p> <p><b>Eared Grebe</b> (<i>Podiceps nigricollis</i>)</p> <p>Franklin's Gull (<i>Larus pipixcan</i>)</p> <p><b>Forster's Tern</b> (<i>Sterna forsteri</i>)</p> <p>Gadwall (<i>Anas strepera</i>)</p> <p><b>Great Blue Heron</b> (<i>Ardea herodias</i>)</p> <p>Hooded Merganser (<i>Lophodytes cucullatus</i>)</p>	<p><b>Horned Lark</b> (<i>Eremophila alpestris</i>)</p> <p><b>Killdeer</b> (<i>Charadrius vociferous</i>)</p> <p><b>Lesser Scaup</b> (<i>Aythya affinis</i>)</p> <p><b>Long-billed Dowitcher</b> (<i>Limnodromus scolopaceus</i>)</p> <p><b>Mallard</b> (<i>Anas platyrhynchos</i>)</p> <p><b>Marsh Wren</b> (<i>Cistothorus palustris</i>)</p> <p><b>Northern Harrier</b> (<i>Circus cyaneus</i>)</p> <p><b>Northern Pintail</b> (<i>Anas acuta</i>)</p> <p>Northern Rough-winged Swallow (<i>Stelgidopteryx serripennis</i>)</p> <p><b>Northern Shoveler</b> (<i>Anas clypeata</i>)</p> <p><b>Raven</b> (<i>Corvus corax</i>)</p> <p>Plover (<i>Charadrius</i> spp.)</p> <p>Red-head Duck (<i>Aythya americana</i>)</p> <p><b>Red-tail Hawk</b> (<i>Buteo jamaicensis</i>)</p> <p><b>Red-winged Blackbird</b> (<i>Agelaius phoeniceus</i>)</p> <p><b>Sandhill Crane</b> (<i>Grus canadensis</i>)</p> <p>Sora (<i>Porzana carolina</i>)</p> <p><b>Tree Swallow</b> (<i>Tachycineta bicolor</i>)</p> <p>Trumpeter Swan (<i>Cygnus buccinator</i>)</p> <p>Tundra Swan (<i>Cygnus columbianus</i>)</p> <p>Vesper Sparrow (<i>Poocetes gramineus</i>)</p> <p>Western Bluebird (<i>Sialia mexicana</i>)</p> <p><b>Western Meadowlark</b> (<i>Sturnella neglecta</i>)</p> <p><b>Wilson's Phalarope</b> (<i>Phalaropus tricolor</i>)</p> <p><b>Yellow-headed Blackbird</b> (<i>Xanthocephalus xanthocephalus</i>)</p>
<b>MAMMALS</b>	
<p><b>Coyote</b> (<i>Canis latrans</i>)</p> <p>Mule Deer (<i>Odocoileus hemionus</i>)</p> <p><b>Muskrat</b> (<i>Ondatra zibethicus</i>)</p> <p>Northern River Otter (<i>Lutra canadensis</i>)</p>	<p>Red Fox (<i>Vulpes vulpes</i>)</p> <p>Striped Skunk (<i>Mephitis mephitis</i>)</p> <p><b>White-tailed Deer</b> (<i>Odocoileus virginianus</i>)</p>

<sup>1</sup>**Bolded** species indicate those observed during 2006.

In 2006 there were almost exactly the same number of birds and bird species observed as in 2005. Slight changes may be due to the specific times and dates observations were made. The greatest number of birds observed at the site in 2006 was 453 compared with over 500 in 2001 and about 200 in 2003. Specific evidence observed, as well as activity codes pertaining to birds, is provided on the completed monitoring form in **Appendix B**.

This site provides habitat for a variety of wildlife species. Three mammal and 29 bird species were noted at the mitigation site during the 2006 site visits (**Table 5**). **Appendix D** includes a list of 81 bird species observed at the site by MDT biologists in recent years.

### 3.6 Macroinvertebrates

Complete results from the four macroinvertebrate sampling collections are presented in **Appendix F**. Samples were not taken at locations 2 and 4 in 2006 due to a lack of water. The following analysis and **Chart 5** were provided by Rhithron Associates, Inc. (Bollman 2006).


**Beaverhead #1.** *Invertebrate abundance appears to have decreased precipitately between 2005 and 2006; only 21 organisms were present in the sample collected this year. Bioassessment is not reliable when too few organisms are collected. Assuming adequate sampling effort, 2006 assessment results suggest diminished water quality and loss of habitat complexity since the earlier year. This follows 5 years of apparent stability in invertebrate assemblage composition and function. Interestingly, proportional representation of functional groups and taxonomic components remained similar between the years, suggesting that the site may have been undersampled in 2006. While sub-optimal biotic conditions were indicated in 2005, scores in 2006 indicated poor conditions.*

**Beaverhead #3.** *Assessment scores remained steady between 2005 and 2006; however, taxa richness increased, and the overall assemblage sensitivity also increased slightly. The invertebrate community composition suggests that open water habitats and habitats dominated by vegetation contributed about equally to complexity at this site. Evidence for the presence of filamentous algae, which was not strong in 2005, appears in 2006 in the large numbers of midges in *Cricotopus (Isocladius) spp.* Dominance by microcrustacea (67% of sampled organisms in 2006) suggests that the site may suffer dewatering periodically. Scores indicated that biotic condition remained poor.*

**Beaverhead #5.** *In 2005, moderate richness of invertebrate taxa was accompanied by low abundance in the sample collected at this site. In 2006 however, very low taxa richness (4) and high abundance characterized the sample collected here. The sampled assemblage was overwhelmed by ostracods. Poor conditions are indicated; aquatic habitats were probably limited. Periodic dewatering or thermal stress cannot be ruled out.*

**Beaverhead #6.** *An assemblage dominated by snails in 2005 shifted to one dominated by mayflies (mainly *Caenis sp.*) and midges (especially *Dicrotendipes sp.* and *Paratanytarsus sp.*) by 2006. The invertebrate fauna collected at this site in 2005 suggests that macrophyte surfaces provided a large component of available habitats. Invertebrate taxa richness was relatively high and relatively stable over the studied years here. Assessment scores indicated sub-optimal conditions.*

**Chart 5: Bioassessment scores for Beaverhead Gateway Wetland Mitigation Site from 2001 to 2006.**



### 3.7 Functional Assessment

The functional assessment numbers for 2006 are similar to those from past years, although a slightly higher recreation/education score was afforded in 2004-2006 as the landowner clarified that permission has and can be granted for birding and scientific research. Further, an interpretive observation area has been constructed overlooking the site along Highway 41. A completed functional assessment form is included in **Appendix B**. The Beaverhead Gateway mitigation site is currently rated as a Category II (high value) site, primarily due to high wildlife habitat, TE habitat, MNHP species habitat, surface water storage, sediment/nutrient removal, food chain support and groundwater discharge ratings. The site received a moderate fish habitat rating due to few fish and habitat deficiencies. The site received a low rating for sediment/shoreline stability due to a lack of plants with deep binding roots, especially along the dike where new fill was placed in 2004 and vegetation is just establishing.

Based on functional assessment results (**Table 6**), approximately 972 functional units have been created thus far at the Beaverhead Gateway mitigation site.

**Table 6: Summary of 2006 wetland function/value ratings and functional points for the Beaverhead Gateway Wetland Mitigation Site.**

Function and Value Parameters From the 1999 MDT Montana Wetland Assessment Method	2006
Listed/Proposed T&E Species Habitat	Mod (0.7)
MTNHP Species Habitat	High (1.0)
General Wildlife Habitat	High (0.9)
General Fish/Aquatic Habitat	Mod (0.5)
Flood Attenuation	Mod (0.5)
Short and Long Term Surface Water Storage	High (1.0)
Sediment, Nutrient, Toxicant Removal	High (1.0)
Sediment/Shoreline Stabilization	Low (0.3)
Production Export/Food Chain Support	High (0.9)
Groundwater Discharge/Recharge	High (1.0)
Uniqueness	Low (0.3)
Recreation/Education Potential	Mod (0.5)
Actual Points/Possible Points	8.6 / 12.0
% of Possible Score Achieved	72%
Overall Category	II
<b>Total Acreage of Assessed Wetlands and Other Aquatic Habitats</b>	<b>118.2</b>
<b>Functional Units (acreage x actual points)</b>	<b>1016.52</b>
<b>Net Acreage Gain</b>	<b>113</b>
<b>Net Functional Unit Gain</b>	<b>971.8</b>

### 3.8 Photographs

Representative photographs taken from photo-points and transect ends are presented in **Appendix C**. A 2001-2006 aerial photograph comparison was compiled (**Appendix C**).

### 3.9 Maintenance Needs/Recommendations

Weed control and revegetation of disturbed sites is still needed to prevent further weed spread, reduce the risk of new weeds invading, reduce wind and water erosion and reduce sediment input to surface waters. Several noxious weeds are present including Canada thistle, hound’s-tongue, and spotted knapweed.

Spoil piles left from ditch excavation continue to create a weed problem, a wind and water erosion hazard and a sedimentation source. This same issue applies to portions of the dike and other poorly vegetated sites. A possible remedy would entail chemically treating weeds and re-seeding the spoil piles with desirable grasses.

Dike erosion and sediment production from the poorly vegetated shoreline could be monitored more closely by installing permanent markers or by periodic surveys. MDT was monitoring erosion on the dike using bank pins from 1998-2001, but the pins are no longer present indicating that erosion has occurred (Urban pers. comm.). The dike was examined by a DNRC

dam inspector in 2005, and the erosion was considered to be a maintenance issue, but not a dam safety concern (see letter in **Appendix D**). The DNRC recommended periodically adding fill to the face of the dike where breaching is taking place and vertical slopes are developing. Fill was added to the face of the dike in 2004 to replace eroded material, but has not yet vegetated. Additional examples of potential solutions to erosion problems include shoreline reinforcement, off-shore wave protection, protected off-shore plantings, shoreline plantings, and placement of vegetated sod mats.

### **3.10 Current Credit Summary**

As of 2006, approximately 118.2 acres of wetland and open water occur within the monitoring area. Subtracting the pre-existing 5.2 wetland acres yields a net of 113 acres, of which 106.5 acres is wetland and 6.5 acres is open water. This includes portions of the monitoring area both above (net of 86.2 wetland acres and 6.5 open water acres) and below (20.3 wetland acres) the dike. MDT has opted not to purchase the credits that have developed below the dike (Urban pers. comm.).

Consequently, available credit at the site (above the dike) as of 2006 is 92.7 aquatic habitat acres, well in excess of the original 52-acre goal. Aquatic habitat features have remained constant in size at the site over the past six years and seem unlikely to change significantly in future years.

#### 4.0 REFERENCES

- Berglund, J. 1999. *MDT Montana Wetland Assessment Method*. May 25<sup>th</sup>. Prepared for Montana Department of Transportation and Morrison-Maierle, Inc. Prepared by Western EcoTech. Helena, Montana. 18 pp.
- Bollman, W. 2006. MDT Mitigated Wetland Monitoring Project – Aquatic Invertebrate Monitoring Summary 2001-2006. Rhithron Associates Inc. Missoula, MT.
- Carlson, J. 2001. Program Zoologist, Montana Natural Heritage Program, Helena, Montana. April conversation.
- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Delineation Manual*. U.S. Army Corps of Engineers, Washington, DC.
- Hackley, Pam. 1997. Pre-Project Wetland Delineation – Beaverhead Gateway Wetland Mitigation Site. Helena, Montana.
- Ralph, C.J., Geupel, G.R., Pyle, P., Martin, T.E., and D.F. DeSante. 1993. *Handbook of field methods for monitoring landbirds*. Gen. Tech. Rep. PSW-GTR-144. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Dept. of Agriculture. 41 p.
- Urban, L. 2001. Wetland Mitigation Specialist, Montana Department of Transportation. Helena, Montana. March 2001 meeting.
- Urban, L. 2002. Wetland Mitigation Specialist, Montana Department of Transportation. Helena, Montana. January 2002 meeting.
- Urban, L. 2004. Wetland Mitigation Specialist, Montana Department of Transportation. Helena, Montana. March 2004 and June 2004 meetings.
- USDA Natural Resource Conservation Service (NRCS). 1989. *Soil Survey of Madison County Area, Montana*.
- USDA Natural Resources Conservation Service. 1998. *Field Indicators of Hydric Soils in the United States*, Version 4. G. Hurt, P. Whited and R. Pringle (eds.). USDA, NRCS Fort Worth, Texas.

## **Appendix A**

---

### **FIGURES 2 & 3**


---

*MDT Wetland Mitigation Monitoring  
Beaverhead Gateway  
Dillon, Montana*

# Figure 2 - 2006 Monitoring Activity Locations

**LEGEND**

- Photograph Point
  - Observation Point
  - Vegetation Transect
  - △ Aerial Reference Point
  - ⊙ Macro-invertebrate Sample Point
- Base Photograph Date: July 14, 2006




PROJECT NAME <b>MDT Beaverhead Gateway Ranch Wetland Mitigation</b>	
DRAWING TITLE <b>2006 Monitoring Activity Locations</b>	
PROJ NO: B43045.011	DRAWN: LLL
LOCATION: Beaverhead Rock	PROJ MGR: J. Berglund
SCALE: 1"=300'	CHECKED: B.D. APP'VD: J.B.
FILE NAME: L:\330054.202\Beaverhead\dwg\Beaverhead2006.dwg	
1120 Cedar Missoula, MT 59802	
<b>PBS</b>	
FIGURE <b>2</b> OF 3	
REV - Dec/11/2006	

# Figure 3 - 2006 Mapped Site Features

- Vegetation Communities:**
- ① Open Water
  - ② Scirpus
  - ③ Hordeum/Kochia-Upland
  - ④ Muhlenbergia/Agropyron-Upland
  - ⑤ Alopecurus/Juncus
  - ⑥ Alopecurus/Scirpus
  - ⑦ Sarcobatus/Elymus-Upland
  - ⑧ Potamogeton/Polygonum-Aquatic

- LEGEND**
- Monitoring Area Limits
  - Open Water Boundary
  - Wetland - Upland Boundary
  - Vegetation Community Boundary
- Base Photograph Date: July 14, 2006

	Monitoring Area	Above Dike	Below Dike
Gross Wetland Area	118.2 Acres	97.9 Acres	20.3 Acres
Open Water Area	6.5 Acres	6.5 Acres	0.0 Acres
Net Wetland Area	111.7 Acres	91.4 Acres	20.3 Acres



PROJECT NAME	MDT Beaverhead Ranch Wetland Mitigation		
DRAWING TITLE	2006 Mapped Site Features		
PROJ NO:	B43054.202	DRAWN:	LLL
LOCATION:	1120 Cedar Missoula, MT 59802	PROJ MGR:	J. Berglund
SCALE:	1" = 300'	CHECKED:	B.D. APP'VD: J.B.
FILE NAME:	L:\330054.202\Beaverhead\dwg\Beaverhead2006.dwg		

## **Appendix B**

---

**2006 WETLAND MITIGATION SITE MONITORING FORM**  
**2006 BIRD SURVEY FORM**  
**2006 COE WETLAND DELINEATION FORMS**  
**2006 MDT FUNCTIONAL ASSESSMENT FORM**

---

*MDT Wetland Mitigation Monitoring*  
*Beaverhead Gateway*  
*Dillon, Montana*

# PBS&J / MDT WETLAND MITIGATION SITE MONITORING FORM

Project Name: Beaverhead Rock Project Number: B43054.00-0202 Assessment Date: 8/9/06  
 Location: NE of Dillon MDT District: Butte Milepost: \_\_\_\_\_  
 Legal description: T 5S R 7W Section 21, 27, & 28 Time of Day: All  
 Weather Conditions: Cloudy Person(s) conducting the assessment: B. Dutton  
 Initial Evaluation Date: \_\_\_/\_\_\_/\_\_\_ Visit #: 6 Monitoring Year: 2006  
 Size of evaluation area: 147 acres Land use surrounding wetland: Agriculture (crops & grazing)

Monitoring area includes wetland & upland.

## HYDROLOGY

**Surface Water** Source: \_\_\_\_\_

Inundation: Present  Absent \_\_\_\_\_ Average depths: 0.25 ft Range of depths: 0 - 4 ft

Assessment area under inundation: 5 %

Depth at emergent vegetation-open water boundary: 1.5 ft

If assessment area is not inundated are the soils saturated w/in 12" of surface: Yes \_\_\_\_\_ No \_\_\_\_\_

Other evidence of hydrology on site (drift lines, erosion, stained vegetation etc.): Drift lines, stained vegetation, drainage patterns, and oxidized root channels.

### Groundwater

Monitoring wells: Present \_\_\_\_\_ Absent

Record depth of water below ground surface

Well #	Depth	Well #	Depth	Well #	Depth

### Additional Activities Checklist:

Map emergent vegetation-open water boundary on air photo

Observe extent of surface water during each site visit and look for evidence of past surface water elevations (drift lines, erosion, vegetation staining etc.)

NA GPS survey groundwater monitoring wells locations if present

**COMMENTS/PROBLEMS:** Site is large and variable. It's difficult to group areas into vegetation types that are narrowly defined without having hundreds of small polygons. Vegetation types as mapped have varying coverage of the indicator species.

---



---

## VEGETATION COMMUNITIES

Community No.: 2 Community Title (main species): Scirpus

Dominant Species	% Cover	Dominant Species	% Cover
<i>Scirpus americanus</i>	9		
<i>Scirpus acutus</i>	P		

**COMMENTS/PROBLEMS:** Bulrush along shorelines- also occurs elsewhere than where shown on map but areas are too small to delineate.

NOTE: # 1 is open water on map.

Community No.: 3 Community Title (main species): Hordeum / Kochia

Dominant Species	% Cover	Dominant Species	% Cover
<i>Hordeum jubatum</i>	2	<i>Agropyron trachycaulum</i>	P
<i>Kochia scoparia</i>	2	<i>Distichlis spicata</i>	P
<i>Cirsium arvense</i>	1	<i>Suaeda intermedia</i>	P
<i>Cardaria draba</i>	P	<i>Descurainia sophia</i>	P
<i>Chenopodium album</i>	T		

**COMMENTS/PROBLEMS:** Weedy community on dikes. Species composition varies.

---



---



---

Community No.: 4 Community Title (main species): Muhlenbergia / Agropyron

Dominant Species	% Cover	Dominant Species	% Cover
<i>Muhlenbergia asperifolia</i>	5	<i>Suaeda intermedia</i>	T
<i>Agropyron smithii</i>	2	<i>Sarcobatus vermiculatus</i>	T
<i>Hordeum jubatum</i>	T	<i>Juncus balticus</i>	T
<i>Elymus cinereus</i>	P	<i>Agropyron trachycaulum</i>	P
<i>Poa pratensis</i>	T		

**COMMENTS/PROBLEMS:** Slightly higher mound above wetland area.

---



---



---

**Additional Activities Checklist:**

X Record and map vegetative communities on air photo

## VEGETATION COMMUNITIES (continued)

Community No.: 5 Community Title (main species): Alopecurus / Juncus

Dominant Species	% Cover	Dominant Species	% Cover
<i>Alopecurus pratensis</i>	7	<i>Rumex crispus</i>	P
<i>Triglochin maritima</i>	P	<i>Agropyron trachycaulum</i>	P
<i>Agrostis alba</i>	1	<i>Carex limnophila</i>	T
<i>Carex nebrascensis</i>	1	<i>Muhlenbergia asperifolia</i>	P
<i>Juncus balticus</i>	1		

**COMMENTS/PROBLEMS:** This area is highly variable. It is dominated by these species but their coverage varies across this community type. Variation is in part due to the transition to wetland character.

---



---

Community No.: 6 Community Title (main species): Alopecurus / Scirpus

Dominant Species	% Cover	Dominant Species	% Cover
<i>Alopecurus pratensis</i>	5	<i>Carex limnophila</i>	T
<i>Scirpus americanus</i>	1	<i>Agropyron trachycaulum</i>	T
<i>Scirpus acutus</i>	P	<i>Scirpus pungens</i>	T
<i>Juncus balticus</i>	2	<i>Hordeum jubatum</i>	T
<i>Triglochin maritima</i>	1	<i>Chenopodium album</i>	T

**COMMENTS/PROBLEMS:** This community is also highly variable on a micro-site basis due to small topographic changes and due to increasing wetlands influence.

---

Community No.: 7 Community Title (main species): Sarcobatus / Elymus

Dominant Species	% Cover	Dominant Species	% Cover
<i>Sarcobatus vermiculatus</i>	3	<i>Juncus balticus</i>	T
<i>Elymus cinereus</i>	1	<i>Poa pratensis</i>	T
<i>Hordeum jubatum</i>	1		
<i>Agropyron smithii</i>	P		
<i>Agropyron trachycaulum</i>	1		

**COMMENTS/PROBLEMS:** Upland areas adjacent to wetland. Similar to 2005.

---



---



---

## COMPREHENSIVE VEGETATION LIST

Species	Vegetation Community Number(s)	Species	Vegetation Community Number(s)
<i>Agropyron cristatum</i>	3,4,7	<i>Epilobium palustris</i>	2,5,6
<i>Agropyron repens</i>	3,4	<i>Equisetum laevigatum</i>	2,5,6
<i>Agropyron smithii</i>	3,4,7	<i>Festuca idahoensis</i>	4,7
<i>Agropyron trachycaulum</i>	3,4	<i>Festuca pratensis</i>	3,7
<i>Agrostis stolonifera</i>	3,4,5,6	<i>Gentianella amarelle</i>	5
<i>Alopecurus pratensis</i>	5,6	<i>Glaux maritime</i>	5,6
<i>Artemisia frigida</i>	3,4,7	<i>Grindelia squarrosa</i>	3,7
<i>Artemisia spp.</i>	7	<i>Habenaria dilatata</i>	2,5,6
<i>Aster falcatus</i>	5	<i>Haplopappus carthamoides</i>	5
<i>Aster hesperius</i>	2,6	<i>Helianthus nuttalli</i>	5
<i>Astragalus spp.</i>	4,7	<i>Hippuris vulgaris</i>	2,6
<i>Bromus inermis</i>	3,4,7	<i>Hordeum jubatum</i>	3,4,7
<i>Bromus japonicus</i>	3,4,7	<i>Iris missouriensis</i>	5
<i>Bromus tectorum</i>	3,7	<i>Iva axillaries</i>	5,6
<i>Calamagrostis neglecta</i>	3,4,7	<i>Juncus balticus</i>	4,5,6
<i>Cardaria draba</i>	3,4,7	<i>Juncus bufonius</i>	2,6
<i>Carex capillaries</i>	5,6	<i>Juncus ensifolius</i>	2,6
<i>Carex limnophila</i>	5,6	<i>Kochia scoparia</i>	3,4
<i>Carex nebrascensis</i>	5,6	<i>Lactuca serriola</i>	3,4,7
<i>Carex praegracilis</i>	5,6	<i>Lepidium perfoliatum</i>	5
<i>Carex spp.</i>	2,5,6	<i>Lycopus asper</i>	2,6
<i>Centaurea maculosa</i>	3,7	<i>Medicago lupulina</i>	5
<i>Chenopodium album</i>	5,6	<i>Medicago sativa</i>	3
<i>Chenopodium rubrum</i>	5,6	<i>Melilotus alba</i>	3,7
<i>Chrysothamnus nauseosus</i>	3,7	<i>Melilotus officinalis</i>	3,7
<i>Cirsium arvense</i>	3,5,6	<i>Mentha arvensis</i>	5,6
<i>Cirsium undulatum</i>	5	<i>Mimulus spp.</i>	5,6
<i>Cleome serrulata</i>	5	<i>Muhlenbergia asperifolia</i>	3,4
<i>Cynoglossum officinale</i>	3,4,5	<i>Phalaris arundinacea</i>	3,5,7
<i>Dactylis glomerata</i>	3,7	<i>Phleum pratense</i>	3,5,7
<i>Descurainia sophia</i>	5	<i>Phlox longifolia</i>	5
<i>Distichlis spicata</i>	5,6	<i>Phragmites australis</i>	5
<i>Eleocharis acicularis</i>	2,6	<i>Plantago eriopoda</i>	5,6
<i>Eleocharis pauciflora</i>	2,6	<i>Poa pratensis</i>	3,4,5,6,7
<i>Elymus cinereus</i>	3,4,7	<b>Continued onto next page.</b>	

**COMMENTS/PROBLEMS:** No new species in 2006.

---



---



---



---





## PHOTOGRAPHS

Using a camera with a 50 mm lenses and color film take photographs of the following permanent reference points listed in the checklist below. Record the direction of the photograph using a compass. (The first time at each site establish a permanent reference point by setting a ½ inch rebar or fencepost extending 2-3' above ground, survey the location with a resource grade GPS and mark the location on the air photo.)

Checklist:

- One photo for each of the 4 cardinal directions surrounding wetland
- At least one photo showing upland use surrounding wetland – if more than one upland use exists, take additional photos
- At least one photo showing buffer surrounding wetland
- One photo from each end of vegetation transect showing transect

Location	Photo Frame #	Photograph Description	Compass Reading
1		Looking NE along fence and W. across mitigation site.	120 & 300
2		Panoramic looking from SW to NE.	270 – 45
3		Looking NE, emergent vegetation / open water and SW along transect.	45 & 225
4		Looking NE, upland vegetation.	45
5		Looking NE across site.	45
7		Looking E. along pond bank and N. along Transect # 2.	90 & 35
8		Looking S. along Transect # 2.	180
9		Looking SE along pond bank & W. along other bank.	150 & 270
10		Looking NE along spoil pile, weedy community.	45

**COMMENTS/PROBLEMS:** Photo Point 7 did not come out

---



---



---

## GPS SURVEYING

Using a resource grade GPS survey the items on the checklist below. Collect at least 3 location points with the GPS unit set at 5 second recording rate. Record file numbers fore site in designated GPS field notebook

Checklist:

- Jurisdictional wetland boundary
- 4-6 landmarks recognizable on the air photo
- Start and end points of vegetation transect(s)
- Photo reference points
- Groundwater monitoring well locations

**COMMENTS/PROBLEMS:** \_\_\_\_\_

---



---



---

**WETLAND DELINEATION**

(Attach Corps of Engineers delineation forms)

At each site conduct the items on the checklist below:

- Delineate wetlands according to the 1987 Army Corps manual.
- Delineate wetland-upland boundary on the air photo
- Survey wetland-upland boundary with a resource grade GPS survey

**COMMENTS/PROBLEMS:** Similar to 2005 and other past years.

---



---



---



---

**FUNCTIONAL ASSESSMENT**

(Complete and attach full MDT Montana Wetland Assessment Method field forms; also attach abbreviated field forms, if used)

**COMMENTS/PROBLEMS:** \_\_\_\_\_

---



---



---



---

**MAINTENANCE**

Were man-made nesting structures installed at this site? YES  NO \_\_\_\_\_

If yes, do they need to be repaired? YES \_\_\_\_\_ NO

If yes, describe problems below and indicate if any actions were taken to remedy the problems.

Were man-made structures build or installed to impound water or control water flow into or out of the wetland?

YES  NO \_\_\_\_\_

If yes, are the structures working properly and in good working order? YES  NO \_\_\_\_\_

If no, describe the problems below.

**COMMENTS/PROBLEMS:** Lots of weeds along excavation piles.

---



---



---



---



---



---



---



---



---



---









**BIRD SURVEY – FIELD DATA SHEET**

Page 1 of 1

Date: 6/1/06

Survey Time: 8:00-10:30

**SITE:** Beaverhead Ranch (Spring)

Bird Species	#	Behavior	Habitat	Bird Species	#	Behavior	Habitat
American coot	4	F	OW				
bald eagle	1	FO					
blue-winged teal	1	F	OW, MA				
Canada goose	76	F, N	OW, MA				
cinnamon teal	14	F	OW, MA				
cliff swallow	60	F	OW, MA				
forster's tern	1	F, L	MF, OW				
killdeer	18	F, N	UP				
lesser scaup	1	F	OW				
mallard	40	F	OW, MA				
marsh wren	2	F	MA				
northern harrier	8	F	UP, MA				
northern pintail	22	F	OW, MA				
northern shoveler	20	F	OW, MA				
pelican	30	L	OW				
raven	2	F	MA				
red-winged blackbird	8	F, N	MA				
sandhill crane	20	F	MA				
tree swallow	70	F	OW, MA				
western meadowlark	2	F	UP				
Wilson's phalarope	36	F	OW, MA				
yellow-headed blackbird	14	F, N	MA				
<b>Total Birds</b>	<b>450</b>						

<b>Notes:</b>
Partly cloudy
tree swallows are using bluebird nest boxes
One coyote and tracks; deer tracks and pellets
Muskrat tracks and burrow

**Behavior:** BP – one of a breeding pair; BD – breeding display; F – foraging; FO – flyover; L – loafing; N – nesting

**Habitat:** AB – aquatic bed; FO – forested; I – island; MA – marsh; MF – mud flat; OW – open water; SS – scrub/shrub; UP – upland buffer; WM – wet meadow, US – unconsolidated shoreline





**DATA FORM**  
**ROUTINE WETLAND DETERMINATION**  
(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Beaverhead Rock</u> Applicant/Owner: <u>MDT</u> Investigator: <u>B. Dutton</u>	Date: <u>08/09/06</u> County: <u>Beaverhead</u> State: <u>MT</u>
Do Normal Circumstances exist on the site: <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No Is the site significantly disturbed (Atypical Situation)? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No Is the area a potential Problem Area?: <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No (If needed, explain on reverse.)	Community ID: _____ Transect ID: <u>T2</u> Plot ID: <u>1</u>

**VEGETATION**

	Dominant Plant Species	Stratum	Indicator		Dominant Plant Species	Stratum	Indicator
1	<i>Alopecurus pratensis</i>	H	FACW	9			
2	<i>Agropyron trachycaulum</i>	H	FAC	10			
3	<i>Juncus balticus</i>	H	FACW+	11			
4	<i>Carex nebrascensis</i>	H	OBL	12			
5	<i>Rumex crispus*</i>	H	FACW	13			
6	<i>Habenaria dilatata</i>	H	OBL	14			
7				15			
8				16			

Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-). 6/6 = 100%

Hydrophytic vegetation present, wetland plants.

**HYDROLOGY**

Recorded Data (Describe in Remarks): <input type="checkbox"/> Stream, Lake, or Tide Gauge <input type="checkbox"/> Aerial Photographs <input type="checkbox"/> Other <input checked="" type="checkbox"/> No Recorded Data Available	Wetland Hydrology Indicators: <b>Primary Indicators:</b> <input type="checkbox"/> Inundated <input type="checkbox"/> Saturated in Upper 12 Inches <input type="checkbox"/> Water Marks <input type="checkbox"/> Drift Lines <input type="checkbox"/> Sediment Deposits <input type="checkbox"/> Drainage Patterns in Wetlands <b>Secondary Indicators (2 or more required):</b> <input checked="" type="checkbox"/> Oxidized Root Channels in Upper 12 Inches <input type="checkbox"/> Water-Stained Leaves <input type="checkbox"/> Local Soil Survey Data <input checked="" type="checkbox"/> FAC-Neutral Test <input type="checkbox"/> Other (Explain in Remarks)
Field Observations:  Depth of Surface Water: _____ (in.)  Depth to Free Water in Pit: <u>&gt;18</u> (in.)  Depth to Saturated Soil: <u>&gt;18</u> (in.)	
Remarks: Dry year.	

# SOILS

Map Unit Name		Neen silty clay loam		Drainage Class: <u>Somewhat poorly</u>	
(Series and Phase):				Field Observations	
Taxonomy (Subgroup):		<u>Aquic calciorthids</u>		Confirm Mapped Type? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	
<b>Profile Description:</b>					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 – 2	O	10YR 4/2	-	-	Silt loam
2 – 12	A1	10 YR 2/0	-	-	Silt loam
12 – 18+	B2	10 YR 1/1	10 YR 6/6	Few/Faint	Very fine sandy loam
<b>Hydric Soil Indicators:</b>					
<input type="checkbox"/> Histosol		<input type="checkbox"/> Concretions			
<input checked="" type="checkbox"/> Histic Epipedon		<input type="checkbox"/> High Organic Content in surface Layer in Sandy Soils			
<input checked="" type="checkbox"/> Sulfidic Odor		<input type="checkbox"/> Organic Streaking in Sandy Soils			
<input type="checkbox"/> Aquic Moisture Regime		<input type="checkbox"/> Listed on Local Hydric Soils List			
<input checked="" type="checkbox"/> Reducing Conditions		<input type="checkbox"/> Listed on National Hydric Soils List			
<input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors		<input type="checkbox"/> Other (Explain in Remarks)			
Mucky mineral surface soil.					

## WETLAND DETERMINATION

Hydrophytic Vegetation Present?	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	Is this Sampling Point Within a Wetland? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
Wetland Hydrology Present?	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	
Hydric Soils Present?	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	
Remarks:			
Same conditions in past years.			

Approved by HQUSACE 2/92

**DATA FORM**  
**ROUTINE WETLAND DETERMINATION**  
(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Beaverhead Rock</u> Applicant/Owner: <u>MDT</u> Investigator: <u>B. Dutton</u>	Date: <u>08/09/06</u> County: <u>Beaverhead</u> State: <u>MT</u>
Do Normal Circumstances exist on the site: <u>X</u> Yes <u>    </u> No Is the site significantly disturbed (Atypical Situation)? <u>    </u> Yes <u>X</u> No Is the area a potential Problem Area?: <u>    </u> Yes <u>X</u> No (If needed, explain on reverse.)	Community ID: <u>                    </u> Transect ID: <u>T2</u> Plot ID: <u>2</u>

**VEGETATION**

Dominant Plant Species	Stratum	Indicator	Dominant Plant Species	Stratum	Indicator
1 <u>Agropyron trachycaulum</u>	<u>H</u>	<u>FAC</u>	9 <u>Elymus cinereus</u>	<u>H</u>	<u>FACU</u>
2 <u>Muhlenbergia asperifolia</u>	<u>H</u>	<u>FACW</u>	10		
3 <u>Festuca idahoensis</u>	<u>H</u>	<u>FACU</u>	11		
4 <u>Rumex crispus</u>	<u>H</u>	<u>FACW</u>	12		
5 <u>Agropyron smithii</u>	<u>H</u>	<u>FACU</u>	13		
6 <u>Hordeum jubatum</u>	<u>H</u>	<u>FAC+</u>	14		
7 <u>Juncus balticus</u>	<u>H</u>	<u>FACW+</u>	15		
8 <u>Poa pratensis</u>	<u>H</u>	<u>FACU+</u>	16		
Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-).			<u>5/9 = 55%</u>		

**HYDROLOGY**

<p>Recorded Data (Describe in Remarks):</p> <p style="padding-left: 20px;"><u>    </u> Stream, Lake, or Tide Gauge</p> <p style="padding-left: 20px;"><u>    </u> Aerial Photographs</p> <p style="padding-left: 20px;"><u>    </u> Other</p> <p><u>  x  </u> No Recorded Data Available</p> <hr/> <p>Field Observations:</p> <p style="padding-left: 20px;">Depth of Surface Water: <u>          </u> (in.)</p> <p style="padding-left: 20px;">Depth to Free Water in Pit: <u>  &gt;20  </u> (in.)</p> <p style="padding-left: 20px;">Depth to Saturated Soil: <u>  &gt;20  </u> (in.)</p>	<p>Wetland Hydrology Indicators:</p> <p>Primary Indicators:</p> <p><u>    </u> Inundated</p> <p><u>    </u> Saturated in Upper 12 Inches</p> <p><u>    </u> Water Marks</p> <p><u>    </u> Drift Lines</p> <p><u>    </u> Sediment Deposits</p> <p><u>    </u> Drainage Patterns in Wetlands</p> <p>Secondary Indicators (2 or more required):</p> <p><u>    </u> Oxidized Root Channels in Upper 12 Inches</p> <p><u>    </u> Water-Stained Leaves</p> <p><u>    </u> Local Soil Survey Data</p> <p><u>    </u> FAC-Neutral Test</p> <p><u>    </u> Other (Explain in Remarks)</p>
Remarks:  Dry year, no obvious hydrologic indicators.	

# SOILS

Map Unit Name		Neen silty clay loam		Drainage Class: <u>  </u> somewhat poorly	
(Series and Phase):				Field Observations	
Taxonomy (Subgroup):		<u>  </u> Aquic calciorthids		Confirm Mapped Type? <u>  </u> Yes <u>  X  </u> No	
<b>Profile Description:</b>					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 - 4	A	10 YR 3/2	-	-	Silt loam
4 - 8	B1	10 YR 4/3	-	-	Silt loam
8 - 20	B2	10 YR 5/3	-	-	Silt loam
<b>Hydric Soil Indicators:</b>					
<u>  </u> Histosol		<u>  </u> Concretions		<u>  </u> High Organic Content in surface Layer in Sandy Soils	
<u>  </u> Histic Epipedon		<u>  </u> Organic Streaking in Sandy Soils		<u>  </u> Listed on Local Hydric Soils List	
<u>  </u> Sulfidic Odor		<u>  </u> Listed on National Hydric Soils List		<u>  </u> Other (Explain in Remarks)	
<u>  </u> Aquic Moisture Regime					
<u>  </u> Reducing Conditions					
<u>  </u> Gleyed or Low-Chroma Colors					
Upland soil colors and features.					

## WETLAND DETERMINATION

Hydrophytic Vegetation Present? <u>  X  </u> Yes <u>  </u> No	Is this Sampling Point Within a Wetland? <u>  </u> Yes <u>  X  </u> No
Wetland Hydrology Present? <u>  </u> Yes <u>  X  </u> No	
Hydric Soils Present? <u>  </u> Yes <u>  X  </u> No	
Remarks:	
Upland site, same conditions in past years.	

**DATA FORM**  
**ROUTINE WETLAND DETERMINATION**  
(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Beaverhead Rock</u>	Date: <u>08/09/06</u>
Applicant/Owner: <u>MDT</u>	County: <u>Beaverhead</u>
Investigator: <u>B. Dutton</u>	State: <u>MT</u>
Do Normal Circumstances exist on the site: <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Community ID: _____
Is the site significantly disturbed (Atypical Situation)? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Transect ID: <u>T1</u>
Is the area a potential Problem Area?: <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No (If needed, explain on reverse.)	Plot ID: <u>3</u>

**VEGETATION**

Dominant Plant Species	Stratum	Indicator	Dominant Plant Species	Stratum	Indicator
1 <u>Sarcobatus vermiculatus</u>	<u>S</u>	<u>FACU+</u>	9 _____		
2 <u>Elymus cinereus</u>	<u>H</u>	<u>FACU</u>	10 _____		
3 <u>Poa pratensis</u>	<u>H</u>	<u>FACU+</u>	11 _____		
4 <u>Agropyron trachycaulum</u>	<u>H</u>	<u>FAC</u>	12 _____		
5 <u>Juncus balticus</u>	<u>H</u>	<u>FACW+</u>	13 _____		
6 _____			14 _____		
7 _____			15 _____		
8 _____			16 _____		
Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-).			<u>2/5 = 40%</u>		
Upland vegetation.					

**HYDROLOGY**

<p>Recorded Data (Describe in Remarks):</p> <p><input type="checkbox"/> Stream, Lake, or Tide Gauge</p> <p><input type="checkbox"/> Aerial Photographs</p> <p><input type="checkbox"/> Other</p> <p><input checked="" type="checkbox"/> No Recorded Data Available</p> <hr/> <p>Field Observations:</p> <p>Depth of Surface Water: _____ (in.)</p> <p>Depth to Free Water in Pit: <u>&gt;18</u> (in.)</p> <p>Depth to Saturated Soil: <u>&gt;18</u> (in.)</p>	<p>Wetland Hydrology Indicators:</p> <p>Primary Indicators:</p> <p><input type="checkbox"/> Inundated</p> <p><input type="checkbox"/> Saturated in Upper 12 Inches</p> <p><input type="checkbox"/> Water Marks</p> <p><input type="checkbox"/> Drift Lines</p> <p><input type="checkbox"/> Sediment Deposits</p> <p><input type="checkbox"/> Drainage Patterns in Wetlands</p> <p>Secondary Indicators (2 or more required):</p> <p><input type="checkbox"/> Oxidized Root Channels in Upper 12 Inches</p> <p><input type="checkbox"/> Water-Stained Leaves</p> <p><input type="checkbox"/> Local Soil Survey Data</p> <p><input type="checkbox"/> FAC-Neutral Test</p> <p><input type="checkbox"/> Other (Explain in Remarks)</p>
<p>Remarks:</p> <p>No hydrologic indicators present.</p>	

# SOILS

Map Unit Name		Neen silty clay loam		Drainage Class: <u>  </u> somewhat poorly	
(Series and Phase):				Field Observations	
Taxonomy (Subgroup):		<u>  </u> Aquic calciorthids		Confirm Mapped Type? <u>  </u> Yes <u>  X  </u> No	
<b>Profile Description:</b>					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 - 7	A1	10 YR 3/2	-	-	Loam
7 - 18	B1	10 YR 4/3	-	-	Loam
<b>Hydric Soil Indicators:</b>					
<u>  </u> Histosol		<u>  </u> Concretions		<u>  </u> High Organic Content in surface Layer in Sandy Soils	
<u>  </u> Histic Epipedon		<u>  </u> Organic Streaking in Sandy Soils		<u>  </u> Listed on Local Hydric Soils List	
<u>  </u> Sulfidic Odor		<u>  </u> Listed on National Hydric Soils List		<u>  </u> Other (Explain in Remarks)	
<u>  </u> Aquic Moisture Regime		<u>  </u> Reducing Conditions			
<u>  </u> Gleyed or Low-Chroma Colors					
Upland soils.					

## WETLAND DETERMINATION

Hydrophytic Vegetation Present?	<u>  </u> Yes	<u>  X  </u> No	Is this Sampling Point Within a Wetland? <u>  </u> Yes <u>  X  </u> No
Wetland Hydrology Present?	<u>  </u> Yes	<u>  X  </u> No	
Hydric Soils Present?	<u>  </u> Yes	<u>  X  </u> No	
Remarks:			
Upland site on small mound above wetland. Same conditions in past years.			



# SOILS

Map Unit Name		Neen silty clay loam		Drainage Class: _____	
(Series and Phase):		_____		Field Observations	
Taxonomy (Subgroup):		<u>Aquic calciorthids</u>		Confirm Mapped Type? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	
<b>Profile Description:</b>					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 - 14	A1	10 YR 2/0	-	-	Loam
14 - 20	B1	10YR 2/1	10 YR 6/6	Few/Faint	Loam
<b>Hydric Soil Indicators:</b>					
<input type="checkbox"/> Histosol		<input type="checkbox"/> Concretions			
<input type="checkbox"/> Histic Epipedon		<input type="checkbox"/> High Organic Content in surface Layer in Sandy Soils			
<input type="checkbox"/> Sulfidic Odor		<input type="checkbox"/> Organic Streaking in Sandy Soils			
<input type="checkbox"/> Aquic Moisture Regime		<input type="checkbox"/> Listed on Local Hydric Soils List			
<input checked="" type="checkbox"/> Reducing Conditions		<input type="checkbox"/> Listed on National Hydric Soils List			
<input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors		<input type="checkbox"/> Other (Explain in Remarks)			
Hydric soil indicators present.					

## WETLAND DETERMINATION

Hydrophytic Vegetation Present?	<u>X</u>	Yes	<input type="checkbox"/>	No	Is this Sampling Point Within a Wetland? <u>x</u> Yes <input type="checkbox"/> No
Wetland Hydrology Present?	<u>X</u>	Yes	<input type="checkbox"/>	No	
Hydric Soils Present?	<u>X</u>	Yes	<input type="checkbox"/>	No	
<b>Remarks:</b>					
Wetland probably will see indicators improve over time as it develops and more natural precipitation levels returns. Same conditions in past years.					

Approved by HQUSACE 2/92

**DATA FORM**  
**ROUTINE WETLAND DETERMINATION**  
(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Beaverhead Rock</u> Applicant/Owner: <u>MDT</u> Investigator: <u>B. Dutton</u>	Date: <u>08/09/06</u> County: <u>Beaverhead</u> State: <u>MT</u>
Do Normal Circumstances exist on the site: <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No Is the site significantly disturbed (Atypical Situation)? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No Is the area a potential Problem Area?: <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No (If needed, explain on reverse.)	Community ID: _____ Transect ID: <u>T1</u> Plot ID: <u>5</u>

**VEGETATION**

Dominant Plant Species	Stratum	Indicator	Dominant Plant Species	Stratum	Indicator
1	<i>Juncus balticus</i>	H	FACW+	9	
2	<i>Spartina gracilis</i>	H	FACW	10	
3	<i>Alopecurus pratensis</i>	H	FACW	11	
4	<i>Chenopodium album</i>	H	FAC	12	
5	<i>Plantago eriopoda</i>	H	FACW	13	
6	<i>Carex limnophila</i>	H	FACW	14	
7	<i>Muhlenbergia asperifolia</i>	H	FACW	15	
8	<i>Agropyron trachycaulum</i>	H	FAC	16	

Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-). 8/8 = 100%

Hydrophytic vegetation present.

**HYDROLOGY**

Recorded Data (Describe in Remarks): <input type="checkbox"/> Stream, Lake, or Tide Gauge <input type="checkbox"/> Aerial Photographs <input type="checkbox"/> Other <input checked="" type="checkbox"/> No Recorded Data Available	Wetland Hydrology Indicators: Primary Indicators: <input type="checkbox"/> Inundated <input type="checkbox"/> Saturated in Upper 12 Inches <input type="checkbox"/> Water Marks <input type="checkbox"/> Drift Lines <input type="checkbox"/> Sediment Deposits <input type="checkbox"/> Drainage Patterns in Wetlands Secondary Indicators (2 or more required): <input checked="" type="checkbox"/> Oxidized Root Channels in Upper 12 Inches <input type="checkbox"/> Water-Stained Leaves <input type="checkbox"/> Local Soil Survey Data <input checked="" type="checkbox"/> FAC-Neutral Test <input type="checkbox"/> Other (Explain in Remarks)
Field Observations: Depth of Surface Water: _____ (in.) Depth to Free Water in Pit: <u>&gt;18</u> (in.) Depth to Saturated Soil: <u>&gt;18</u> (in.)	
Remarks: Dry part of year during multi-year drought cycle. Secondary hydrologic indicators present.	

# SOILS

Map Unit Name		Neen silty clay loam		Drainage Class: _____	
(Series and Phase):		_____		Field Observations	
Taxonomy (Subgroup):		<u>Aquic calciorthids</u>		Confirm Mapped Type? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	
<b>Profile Description:</b>					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 - 2	A1	10 YR 5/4	-		Loam
2 - 18	B1	10 YR 7/1	10 YR 6/6	Few/Faint	Silty clay loam
<b>Hydric Soil Indicators:</b>					
<input type="checkbox"/> Histosol		<input type="checkbox"/> Concretions			
<input type="checkbox"/> Histic Epipedon		<input type="checkbox"/> High Organic Content in surface Layer in Sandy Soils			
<input type="checkbox"/> Sulfidic Odor		<input type="checkbox"/> Organic Streaking in Sandy Soils			
<input checked="" type="checkbox"/> Aquic Moisture Regime		<input type="checkbox"/> Listed on Local Hydric Soils List			
<input type="checkbox"/> Reducing Conditions		<input type="checkbox"/> Listed on National Hydric Soils List			
<input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors		<input type="checkbox"/> Other (Explain in Remarks)			
Soil is developing hydric features, will likely get stronger with more normal rainfall.					

## WETLAND DETERMINATION

Hydrophytic Vegetation Present?	<u>X</u>	Yes	<input type="checkbox"/>	No	Is this Sampling Point Within a Wetland? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
Wetland Hydrology Present?	<u>X</u>	Yes	<input type="checkbox"/>	No	
Hydric Soils Present?	<u>X</u>	Yes	<input type="checkbox"/>	No	
<b>Remarks:</b>					
Soil and hydrology indicators are not very strong, but there, and are likely to improve with normal precipitation. Same conditions in past years.					

**DATA FORM**  
**ROUTINE WETLAND DETERMINATION**  
(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Beaverhead Rock</u> Applicant/Owner: <u>MDT</u> Investigator: <u>B. Dutton</u>	Date: <u>08/09/06</u> County: <u>Beaverhead</u> State: <u>MT</u>
Do Normal Circumstances exist on the site: <u>  x  </u> Yes <u>    </u> No Is the site significantly disturbed (Atypical Situation)? <u>    </u> Yes <u>  X  </u> No Is the area a potential Problem Area?: <u>    </u> Yes <u>  X  </u> No (If needed, explain on reverse.)	Community ID: <u>                    </u> Transect ID: <u>  T1  </u> Plot ID: <u>    6    </u>

**VEGETATION**

Dominant Plant Species	Stratum	Indicator	Dominant Plant Species	Stratum	Indicator
1 <i>Scirpus acutus</i>	H	OBL	9		
2 <i>Hordeum jubatum</i>	H	FAC+	10		
3 <i>Scirpus americanus</i>	H	OBL	11		
4	h		12		
5			13		
6			14		
7			15		
8			16		

Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-).   3/3 = 100%  

Wetland vegetation present.

**HYDROLOGY**

Recorded Data (Describe in Remarks): <u>    </u> Stream, Lake, or Tide Gauge <u>    </u> Aerial Photographs <u>    </u> Other <u>  x  </u> No Recorded Data Available	Wetland Hydrology Indicators: Primary Indicators: <u>    </u> Inundated <u>  x  </u> Saturated in Upper 12 Inches <u>  x  </u> Water Marks <u>  x  </u> Drift Lines <u>    </u> Sediment Deposits <u>    </u> Drainage Patterns in Wetlands Secondary Indicators (2 or more required): <u>    </u> Oxidized Root Channels in Upper 12 Inches <u>    </u> Water-Stained Leaves <u>    </u> Local Soil Survey Data <u>    </u> FAC-Neutral Test <u>    </u> Other (Explain in Remarks)
Field Observations: Depth of Surface Water: <u>            </u> (in.) Depth to Free Water in Pit: <u>    24    </u> (in.) Depth to Saturated Soil: <u>    8    </u> (in.)	
Remarks: Wetland hydrology.	

# SOILS

Map Unit Name		Neen silty clay loam		Drainage Class: _____	
(Series and Phase):		_____		Field Observations	
Taxonomy (Subgroup):		Aquic calciorthids		Confirm Mapped Type? _____ Yes <input checked="" type="checkbox"/> No	
<b>Profile Description:</b>					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 – 2	A1	10 YR 6/3	-	-	Silt loam
2 – 18	B1	10 YR 7/1	10 YR 7/4	-	Loam
<b>Hydric Soil Indicators:</b>					
_____ Histosol		_____ Concretions		_____ High Organic Content in surface Layer in Sandy Soils	
_____ Histic Epipedon		_____ Organic Streaking in Sandy Soils		_____ Listed on Local Hydric Soils List	
_____ Sulfidic Odor		_____ Listed on National Hydric Soils List		_____ Other (Explain in Remarks)	
<input checked="" type="checkbox"/> Aquic Moisture Regime					
_____ Reducing Conditions					
<input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors					
Thin surface layer of more recent deposition over very low chroma and high organic matter layer.					

## WETLAND DETERMINATION

Hydrophytic Vegetation Present? _____ Yes _____ No	Is this Sampling Point Within a Wetland? _____ Yes _____ No
Wetland Hydrology Present? _____ Yes _____ No	
Hydric Soils Present? _____ Yes _____ No	
<b>Remarks:</b>  Good wetland, same conditions in past years.	



**14A. HABITAT FOR FEDERALLY LISTED OR PROPOSED THREATENED OR ENDANGERED PLANTS AND ANIMALS**

i. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species)  D  S \_\_\_\_\_
- Secondary habitat (list species)  D  S Bald Eagle
- Incidental habitat (list species)  D  S \_\_\_\_\_
- No usable habitat  D  S \_\_\_\_\_

ii. Rating (Based on the strongest habitat chosen in 14A(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point & Rating	---	---	---	.7 (M)	---	---	---

If documented, list the source (e.g., observations, records, etc.): \_\_\_\_\_

**14B. HABITAT FOR PLANTS AND ANIMALS RATED AS S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM.**

Do not include species listed in 14A(i).

i. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species)  D  S Black Tern, Lemmons alkaligrass.
- Secondary habitat (list species)  D  S \_\_\_\_\_
- Incidental habitat (list species)  D  S \_\_\_\_\_
- No usable habitat  D  S \_\_\_\_\_

ii. Rating: Based on the strongest habitat chosen in 14B(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point & Rating	1 (H)	---	---	---	---	---	---

If documented, list the source (e.g., observations, records, etc.): Black Terns and Lemmon's alkaligrass documented onsite. Forster's terns and trumpeter swans also observed (but not breeding).

**14C. GENERAL WILDLIFE HABITAT RATING**

i. Evidence of overall wildlife use in the AA: Check either substantial, moderate, or low.

**Substantial** (based on any of the following)

- observations of abundant wildlife #s or high species diversity (during any period)
- abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
- presence of extremely limiting habitat features not available in the surrounding area
- interviews with local biologists with knowledge of the AA

**Low** (based on any of the following)

- few or no wildlife observations during peak use periods
- little to no wildlife sign
- sparse adjacent upland food sources
- interviews with local biologists with knowledge of AA

**Moderate** (based on any of the following)

- observations of scattered wildlife groups or individuals or relatively few species during peak periods
- common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
- adequate adjacent upland food sources
- interviews with local biologists with knowledge of the AA

ii. **Wildlife Habitat Features:** Working from top to bottom, select the AA attribute to determine the exceptional (E), high (H), moderate (M), or low (L) rating. Structural diversity is from 13. For class cover to be considered evenly distributed, vegetated classes must be within 20% of each other in terms of their percent composition in the AA (see 10). Duration of Surface Water: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; A= absent.

Structural Diversity (from 13)	<input type="checkbox"/> High								<input checked="" type="checkbox"/> Moderate								<input type="checkbox"/> Low			
	<input type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input type="checkbox"/> Even				<input checked="" type="checkbox"/> Uneven				<input type="checkbox"/> Even			
	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Class Cover Distribution (all vegetated classes)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Duration of Surface Water in ≥ 10% of AA	--	--	--	--	--	--	--	--	--	--	--	--	H	--	--	--	--	--	--	--
Low disturbance at AA (see 12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (see 12)	--	--	--	--	--	--	--	--	--	--	--	--	H	--	--	--	--	--	--	--
High disturbance at AA (see 12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

iii. Rating: Use 14C(i) and 14C(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L) for this function.

Evidence of Wildlife Use from 14C(i)	Wildlife Habitat Features Rating from 14C(ii)			
	<input type="checkbox"/> Exceptional	<input checked="" type="checkbox"/> High	<input type="checkbox"/> Moderate	<input type="checkbox"/> Low
Substantial	--	.9 (H)	--	--
Moderate	--	--	--	--
Low	--	--	--	--

Comments: Numerous waterfowl and shorebirds observed.

**14D. GENERAL FISH / AQUATIC HABITAT RATING**  NA (proceed to 14E)

If the AA is not or was not historically used by fish due to lack of habitat or excessive gradient, then check the NA box above.

Assess if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [e.g. fish use is precluded by perched culvert or other barrier, etc.]. If fish use occurs in the AA but is not desired from a resource management perspective (e.g. fish use within an irrigation canal), then Habitat Quality [14D(i)] below should be marked as "Low", applied accordingly in 14D(ii) below, and noted in the comments.

**i. Habitat Quality:** Pick the appropriate AA attributes in matrix to determine the quality rating of exceptional (E), high (H), moderate (M), or low (L).

Duration of Surface Water in AA	<input checked="" type="checkbox"/> Permanent/Perennial			<input type="checkbox"/> Seasonal / Intermittent			<input type="checkbox"/> Temporary / Ephemeral		
	>25%	10-25%	<10%	>25%	10-25%	<10%	>25%	10-25%	<10%
Cover - % of waterbody in AA containing cover objects (e.g. submerged logs, large rocks & boulders, overhanging banks, floating-leaved vegetation)									
Shading - >75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities	--	--	--	--	--	--	--	--	--
Shading - 50 to 75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	--	--	--	--	--	--	--
Shading - < 50% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	M	--	--	--	--	--	--

**ii. Modified Habitat Quality:** Is fish use of the AA precluded or significantly reduced by a culvert, dike, other man-made structure or activity or is the waterbody included on the 'MDEQ list of waterbodies in need of TMDL development' with 'Probable Impaired Uses' listed as cold or warm water fishery or aquatic life support?

Y  N If yes, reduce the rating from 14D(i) by one level and check the modified habitat quality rating:  E  H  M  L

**iii. Rating:** Use the conclusions from 14D(i) and 14D(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L).

Types of Fish Known or Suspected within AA	Modified Habitat Quality from 14D(ii)			
	<input type="checkbox"/> Exceptional	<input type="checkbox"/> High	<input type="checkbox"/> Moderate	<input checked="" type="checkbox"/> Low
Native game fish	--	--	--	.5 (M)
Introduced game fish	--	--	--	--
Non-game fish	--	--	--	--
No fish	--	--	--	--

Comments: Unidentified minnows assumed to be native game fish.

**14E. FLOOD ATTENUATION**  NA (proceed to 14F)

Applies only to wetlands subject to flooding via in-channel or overbank flow. If wetlands in AA do not flood from in-channel or overbank flow, then check NA.

**i. Rating:** Working from top to bottom, mark the appropriate attributes to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Estimated wetland area in AA subject to periodic flooding	<input checked="" type="checkbox"/> ≥ 10 acres			<input type="checkbox"/> <10, >2 acres			<input type="checkbox"/> ≤2 acres		
	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
% of flooded wetland classified as forested, scrub/shrub, or both									
AA contains no outlet or restricted outlet	--	--	--	--	--	--	--	--	--
AA contains unrestricted outlet	--	--	.5 (M)	--	--	--	--	--	--

**ii. Are residences, businesses, or other features which may be significantly damaged by floods located within 0.5 miles downstream of the AA? (check)**

Y  N Comments: Potentially flooded area is northeast of dike along river.

**14F. SHORT AND LONG TERM SURFACE WATER STORAGE**  NA (proceed to 14G)

Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow.

If no wetlands in the AA are subject to flooding or ponding, then check NA above.

**i. Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral.

Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding.	<input checked="" type="checkbox"/> >5 acre feet			<input type="checkbox"/> <5, >1 acre feet			<input type="checkbox"/> ≤1 acre foot		
	P/P	S/I	T/E	P/P	S/I	T/E	P/P	S/I	T/E
Duration of surface water at wetlands within the AA									
Wetlands in AA flood or pond ≥ 5 out of 10 years	1 (H)	--	--	--	--	--	--	--	--
Wetlands in AA flood or pond < 5 out of 10 years	--	--	--	--	--	--	--	--	--

Comments: \_\_\_\_\_

**14G. SEDIMENT/NUTRIENT/TOXICANT RETENTION AND REMOVAL**  NA (proceed to 14H)

Applies to wetlands with the potential to receive excess sediments, nutrients, or toxicants through influx of surface or ground water or direct input.

If no wetlands in the AA are subject to such input, check NA above.

**i. Rating** Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Sediment, Nutrient, and Toxicant Input Levels Within AA	AA receives or surrounding land use has potential to deliver low to moderate levels of sediments, nutrients, or compounds such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
	<input checked="" type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%		<input type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%	
% cover of wetland vegetation in AA	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Evidence of flooding or ponding in AA								
AA contains no or restricted outlet	1 (H)	--	--	--	--	--	--	--
AA contains unrestricted outlet	--	--	--	--	--	--	--	--

Comments: Most of the AA has a restricted outlet and is subject to agricultural run-off from cropland to the west.

**14H. SEDIMENT/ShORELINE STABILIZATION**  NA (proceed to 14I)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body that is subject to wave action. If this does not apply, then check NA above.

i. **Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating exceptional (E), high (H), moderate (M), or low (L) for this function.

% Cover of wetland streambank or shoreline by species with deep, binding rootmasses.	Duration of Surface Water Adjacent to Rooted Vegetation		
	<input checked="" type="checkbox"/> Permanent / Perennial	<input type="checkbox"/> Seasonal / Intermittent	<input type="checkbox"/> Temporary / Ephemeral
≥ 65 %	--	--	--
35-64 %	--	--	--
< 35 %	.3 (L)	--	--

Comments: \_\_\_\_\_

**14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT**

i. **Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

A = acreage of vegetated component in the AA. B = structural diversity rating from #13. C = Yes (Y) or No (N) as to whether or not the AA contains a surface or subsurface outlet. P/P = permanent/perennial; S/I = seasonal/intermittent; T/E/A = temporary/ephemeral/absent.

A	<input checked="" type="checkbox"/> Vegetated component >5 acres						<input type="checkbox"/> Vegetated component 1-5 acres						<input type="checkbox"/> Vegetated component <1 acre					
B	<input type="checkbox"/> High		<input checked="" type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low	
C	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input checked="" type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N
P/P	--	--	.9H	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
S/I	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
T/E/A	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Comments: \_\_\_\_\_

**14J. GROUNDWATER DISCHARGE / RECHARGE (DR)** (Check the indicators in i & ii below that apply to the AA.)

i.  **Discharge Indicators**

- Springs are known or observed.
- Vegetation growing during dormant season / drought.
- Wetland occurs at the toe of a natural slope.
- Seeps are present at the wetland edge.
- AA permanently flooded during drought periods.
- Wetland contains an outlet, but no inlet.
- Other \_\_\_\_\_

ii.  **Recharge Indicators**

- Permeable substrate presents without underlying impeding layer.
- Wetland contains inlet but not outlet.
- Other \_\_\_\_\_

iii. **Rating:** Use information from 14J(i) and 14J(ii) above and the table below to arrive at the functional point and rating of high (H) or low (L) for this function.

Criteria	Functional Point and Rating
AA has known Discharge/Recharge area or one or more indicators of D/R present	1 (H)
No Discharge/Recharge indicators present	--
Available Discharge/Recharge information inadequate to rate AA D/R potential	--

Comments: \_\_\_\_\_

**14K. UNIQUENESS**

i. **Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Replacement Potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP.			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP.			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate.		
	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input checked="" type="checkbox"/> common	<input type="checkbox"/> abundant
Estimated Relative Abundance from 11									
Low disturbance at AA (12i)	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (12i)	--	--	--	--	--	--	--	.3L	--
High disturbance at AA (12i)	--	--	--	--	--	--	--	--	--

Comments: \_\_\_\_\_

**14L. RECREATION / EDUCATION POTENTIAL**

i. Is the AA a known recreational or educational site?  Yes [Rate  High (1.0), then proceed to 14L(ii) only]  No [Proceed to 14L(iii)]

ii. Check categories that apply to the AA:  Educational / scientific study  Consumptive rec.  Non-consumptive rec.  Other

iii. Based on the location, diversity, size, and other site attributes, is there a strong potential for recreational or educational use?

- Yes [Proceed to 14L (ii) and then 14L(iv)]
- No [Rate as low in 14L(iv)]

iv. **Rating** Use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Ownership	Disturbance at AA from 12(i)		
	<input type="checkbox"/> Low	<input checked="" type="checkbox"/> Moderate	<input type="checkbox"/> High
Public ownership	--	--	--
Private ownership	--	.3(L)	--

Comments: 0.5 assigned, over-riding calculated score as the landowner will grant permission for scientific study and birding.

**FUNCTION, VALUE SUMMARY, AND OVERALL RATING**

Function and Value Variables	Rating	Actual Functional Points	Possible Functional Points	Functional Units (Actual Points x Estimated AA Acreage)
A. Listed/Proposed T&E Species Habitat	moderate	0.70	1	
B. MT Natural Heritage Program Species Habitat	high	1.00	1	
C. General Wildlife Habitat	high	0.90	1	
D. General Fish/Aquatic Habitat	moderate	0.50	1	
E. Flood Attenuation	moderate	0.50	1	
F. Short and Long Term Surface Water Storage	high	1.00	1	
G. Sediment/Nutrient/Toxicant Removal	high	1.00	1	
H. Sediment/Shoreline Stabilization	low	0.30	1	
I. Production Export/Food Chain Support	high	0.90	1	
J. Groundwater Discharge/Recharge	high	1.00	1	
K. Uniqueness	low	0.30	1	
L. Recreation/Education Potential	moderate	0.50	1	
<b>Total:</b>		<b><u>8.60</u></b>	<b><u>12.00</u></b>	_____
<b>Percent of Total Possible Points:</b>			<b><u>72%</u></b> (Actual / Possible) x 100 [rd to nearest whole #]	

<p><b>Category I Wetland:</b> (Must satisfy <b>one</b> of the following criteria. If not satisfied, proceed to Category II.)</p> <p><input type="checkbox"/> Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; <b>or</b></p> <p><input type="checkbox"/> Score of 1 functional point for Uniqueness; <b>or</b></p> <p><input type="checkbox"/> Score of 1 functional point for Flood Attenuation <b>and</b> answer to Question 14E(ii) is "yes"; <b>or</b></p> <p><input type="checkbox"/> Percent of total Possible Points is &gt; 80%.</p>
<p><b>Category II Wetland:</b> (Criteria for Category I not satisfied <b>and</b> meets any <b>one</b> of the following Category II criteria. If not satisfied, proceed to Category IV.)</p> <p><input checked="" type="checkbox"/> Score of 1 functional point for Species Rated S1, S2, or S3 by the MT Natural Heritage Program; <b>or</b></p> <p><input checked="" type="checkbox"/> Score of .9 or 1 functional point for General Wildlife Habitat; <b>or</b></p> <p><input type="checkbox"/> Score of .9 or 1 functional point for General Fish/Aquatic Habitat; <b>or</b></p> <p><input type="checkbox"/> "High" to "Exceptional" ratings for <b>both</b> General Wildlife Habitat <b>and</b> General Fish / Aquatic Habitat; <b>or</b></p> <p><input type="checkbox"/> Score of .9 functional point for Uniqueness; <b>or</b></p> <p><input checked="" type="checkbox"/> Percent of total possible points is &gt; 65%.</p>
<p><input type="checkbox"/> <b>Category III Wetland:</b> (Criteria for Categories I, II, or IV not satisfied.)</p>
<p><b>Category IV Wetland:</b> (Criteria for Categories I or II are not satisfied <b>and</b> <u>all</u> of the following criteria are met; If not satisfied, return to Category III.)</p> <p><input type="checkbox"/> "Low" rating for Uniqueness; <b>and</b></p> <p><input type="checkbox"/> "Low" rating for Production Export / Food Chain Support; <b>and</b></p> <p><input type="checkbox"/> Percent of total possible points is &lt; 30%.</p>

**OVERALL ANALYSIS AREA (AA) RATING:** (Check appropriate category based on the criteria outlined above.)

- I**
         
  **II**
         
  **III**
         
  **IV**

## **Appendix C**

---

### **2006 REPRESENTATIVE PHOTOGRAPHS 2001-2006 AERIAL PHOTOGRAPH COMPARISON**

---

*MDT Wetland Mitigation Monitoring  
Beaverhead Gateway  
Dillon, Montana*

## 2006 BEAVERHEAD GATEWAY WETLAND MITIGATION SITE



Photo Point No.1: View looking northeast along fence-line (60°).



Photo Point No. 1: View looking northwest across mitigation site. Upland to wetland vegetation transition (300°)



Photo Point No. 3: View looking southwest along the end of Transect 1, emergent wetland vegetation dominated by bulrush (225°).



Photo Point No. 3: View looking northeast, open water and emergent wetland vegetation dominated by bulrush (45°).



Photo Point No. 4: View looking northeast along the beginning of Transect 1 (40°).



Photo Point No. 5: View looking northeast across mitigation site (45°).

**2006 BEAVERHEAD GATEWAY WETLAND MITIGATION SITE**



Photo Point No. 2: Panoramic view of mitigation site, southern half, 300° to 220°. Photo taken looking north to south.



Photo Point No. 2: Panoramic view of mitigation site, northern half, 40° to 300°. Photo taken looking north to south.

**2006 BEAVERHEAD GATEWAY WETLAND MITIGATION SITE**



Photo Point No. 7: View looking north along the start of Transect 2 (350°).



Photo Point No. 8: View looking south from the end of Transect 2 (170°).



Photo Point No. 9: View looking west along dike shore and open water (270°).



Photo Point No. 9: View looking along dike shore and open water (90°).



Photo Point No. 10: View looking northeast along spoil pile dominated by a weedy plant community (45°).

SIX-YEAR AERIAL PHOTOGRAPH COMPARISON – BEAVERHEAD GATEWAY WETLAND MITIGATION SITE




JULY 18, 2001




JULY 22, 2002



JULY 27, 2003



JULY 25, 2004



JULY 27, 2005



JULY 14, 2006


## **Appendix D**

---

**ORIGINAL SITE PLAN  
MDT BIRD OBSERVATIONS  
LETTERS ADDRESSING SITE MANAGEMENT**

---

*MDT Wetland Mitigation Monitoring  
Beaverhead Gateway  
Dillon, Montana*



Beavee head Ranch Mitigation  
Proposed WATER DEPTHS

SYMBOL	DESCRIPTION	ACTUAL ACREAGE	CREDIT ACREAGE
	GROUNDWATER DEPTH FROM 15' TO 24'	18.5 AC.	4.7 AC.
	GROUNDWATER DEPTH FROM 0' TO 12'	22.6 AC.	22.6 AC.
	STANDING WATER DEPTH FROM 0' TO 12'	11.7 AC.	11.7 AC.
	STANDING WATER DEPTH FROM 12' TO 24'	11.3 AC.	11.3 AC.
	STANDING WATER WITH GREATER THAN 24'	2.8 AC.	2.8 AC.
			<b>53.1 AC.</b>

- WATER TEMPERATURE SAMPLING STATIONS
- MACROINVERTEBRATE SAMPLING SITES

**BEAVERHEAD GATEWAY RANCH  
WETLAND MITIGATION SITE  
ACTIVE BIRD LIST 1997 to Present**

(b) breeding

**Waterfowl:**

Tundra Swan  
Trumpeter Swan  
Blue-winged Teal (b)  
Green-winged Teal  
Cinnamon Teal (b)  
Mallard Duck (b)  
Pintail Duck (b)  
Ruddy Duck (b?)  
Greater Canada Geese (b)  
Snow Geese  
Northern Shoveller (b)  
American Wigeon (b)  
Redhead Duck (b)  
Gadwall (b)  
Bufflehead (b)  
Common Goldeneye  
Barrow's Goldeneye  
Lesser Scaup  
American Coot (b)  
Western Grebe  
Eared Grebe (b)  
Double Crested Cormorants  
Red-breasted Merganser  
Common Merganser

**Hérons / Cranes:**

Great Blue Heron  
Black Crowned Night Heron  
Sandhill Cranes (b)

**Eagles / Hawks:**

Golden Eagle  
Red-Tailed Hawk  
Merlin  
American Kestrel  
Northern Harrier  
Rough-legged Hawk  
Peregrine Falcon

**Shorebirds:**

American Avocet  
Willet  
Marbled Godwit  
Wilson's Phalarope  
Red Phalarope (b)  
Common Snipe (b)  
Solitary Sandpiper  
Spotted Sandpiper (b)  
Killdeer (b)  
Greater Yellowlegs  
Sanderlings  
Lesser Yellowlegs  
Long-billed Dowitcher

**Gulls / Terns:**

Franklin's Gull  
Bonaparte's Gull  
Common Tern  
Black Tern (b?)

**Swallows / Swifts:**

Bank Swallows (b)  
Cliff Swallows (b)  
Violet-green Swallows (b)  
Barn Swallows (b)

**Upland Gamebirds:**

Ring-necked Pheasant  
Sage Grouse  
Chukar  
Hungarian Partridge (b)

**Dippers:**

American Dipper

**Owls:**

Short-eared owl

**Crows / Ravens:**

American Crow  
Common Raven  
Black-billed Magpie

**Songbirds:**

Red-winged Blackbird (b)  
Yellow-headed blackbird (b)  
Brewer's Blackbird  
Vesper Sparrow (b)  
Song Sparrow  
Savannah Sparrow (b)  
Western Bluebirds (b)  
American Robin  
American Goldfinch (b)  
Brown-headed Cowbird  
Western Meadowlark (b)  
European Starling  
Mourning Dove  
Rock Dove  
Spotted Towhee

**Pelicans:**

American White Pelican

June 21, 2002

Mr. James T. Harrison, Jr.  
Attorney at Law  
1330 Helena Avenue  
P.O. Box 6876  
Helena, MT 59604-6876

Subject: Beaverhead Gateway Ranch Site Visit of June 17, 2002

Dear Tom:

Joel Marshik, Lyle Manley and I enjoyed meeting with you and Mr. Peccia on June 17 and were encouraged that we are moving in the right direction. We discussed grazing and development of a grazing management plan, Mr. Peccia's concern regarding possible liability for fire passing through the easement property and causing harm to third parties, dike stability and erosion, weed control, the spoil piles, removal of debris, and closure of the on-site well.

I would like to briefly summarize our discussions of the 17<sup>th</sup> and propose a next step. If my recollection of our discussions is incorrect in any way, please advise.

**Grazing and managed grazing strategy:** we will meet on July 8, 2002 to further discuss this issue.

**Fire:** Mr. Peccia expressed concern about the possibility of liability arising from a fire event that passes through the easement area and then on to adjoining property on which damage occurs to property owned by a third party. I perceived that Mr. Peccia was concerned about a fire that was not started by either him (or his employees/agents) or personnel from MDT, but a fire caused by the actions of a third party or an act of God which simply passes through the easement property. It is my opinion that the risk imposed upon Mr. Peccia for liability under this scenario is remote. I am unaware of any court holding which designates a wetland, restored and monitored pursuant to federal law and regulations, a nuisance, or which states that the property owner, in so restoring and maintaining the wetland, could be held liable for any damages to third parties under any theory of law. Given this, and in the interest of continuing negotiations, Lyle Manley and I will investigate whether MDT would be willing to indemnify Mr. Peccia against such a remote risk. Under no circumstances can MDT enter into such an agreement if it could be interpreted that MDT somehow agrees to assume liability for damage to the property of third parties. Lyle and I will contact you after we have further investigated this issue.

**Dike erosion:** Significant erosion of the dike has occurred due to wind and wave action on the highly erodible material of which the dike is construction. In an area near the west end of the dike, vegetation has established. Along the rest of the dike, vegetation appears to be establishing slowly. Mr. Peccia discussed the possibility of utilizing additional rubber mat (terminology?) along the face of the dike and also lowering the water level to expose saturated soil at the eroded area of the dike so that vegetation may establish. As stated by Joel Marshik, additional matting should be installed and it may be possible to lower the water level for further establishment of vegetation and for flushing of the control mechanism. Additionally, due to the fractures along the face of the dike caused by erosion, further slumping of dike material may occur and should be closely monitored. Should further measures become necessary in the future to insure the stability of the dike, MDT is willing to discuss alternatives.

**Weeds:** Several patches of whitetop were noted during the tour. Mr. Peccia indicated that those weeds, and others noted by the Land & Water monitoring report, will be treated.

**Spoil Piles:** I have been advised that my initial statement to you and Mr. Peccia that MDT wanted the spoil piles removed was incorrect and I apologize for my misstatement. I have since learned that it was recommended that the spoil piles remain, but be manipulated to provide accessible habitat and passage for upland wildlife including weed eradication and establishment of native upland vegetation.

**Debris:** As noted during the site visit, the shop area still contains a significant amount of debris resulting from the demolition of the shop, fences and other structures. Mr. Peccia stated that he intends to dispose of the debris.

**Well:** Mr. Peccia stated that he would see to the closure of the well located on the easement area.

When the original agreement was negotiated, MDT was assured that the above-discussed items would be addressed as part of the development of the area. Negotiations on the additional purchase and trade to amend the original easement would be significantly furthered if Mr. Peccia would agree to discuss with MDT a methodology and timetable setting forth dates by which the above-discussed items will be addressed. Such a document should include methods and dates for: 1) installation of further erosion prevention measures on the dike; 2) weed control treatment; 3) habitat restoration and weed control on the spoil piles; 4) removal of debris from the old shop area; and 5) closure of the well. Additionally, as stated above, lowering the level of water to allow further vegetation establishment and flushing of the control structure may be appropriate. Please get back to me on whether Mr. Peccia would be agreeable to address these issues.

Additionally, Lyle Manley and I will get back to you regarding the possibility of a fire indemnification provision, and we look forward to discussing the grazing strategy with you on July 8. Please do not hesitate to contact me if I may be of further service.

Sincerely,

Robert M. Gentry  
Attorney  
MDT Legal Services  
(406) 444-3237

Cc: Joel Marshik  
Gordon Stockstad  
Lyle Manley  
Larry Urban

January 26, 2004

RECEIVED

JAN 27 2004

ENVIRONMENTAL

Bonnie Steg  
Resources Section Supervisor  
Environmental Services Bureau  
Montana Department of Transportation  
2701 Prospect Avenue  
P.O. Box 201001  
Helena, MT 59620-1001

Subject: STPX 0002 (624) CN# 3476  
Beaverhead Gateway Ranch  
U.S. Army Corps of Engineers  
Restrictions

Bonnie Steg,

As I reply to your letter of January 21, 2004 please keep in mind that I am simply trying to state my positions and opinions with clarity. I would appreciate the full context of the Corps letter so that I can be sure to address their concerns, and would also appreciate knowing how many wetlands credits have been withdrawn from the wetland by the Department so I can evaluate how close I am to my contractual requirements. I would also like to know if the Corps has made a personal inspection of the dike or if they have made their determination from reading the Wetland Mitigation Monitoring Report.

You stated in your letter that the erosion on the dike, as indicated in previous letters and meetings, "is a re-occurring problem that needs to be addressed and corrected before the structure fails and repair costs escalate". I have discussed this issue many times before with the Department and I have tried to explain that the material on the front face of the dike is not structural. Until the erosion begins to affect the clay core the integrity of the dike is not compromised.

My objective has always been to create a natural wetland with a natural looking dike using vegetative cover for erosion protection. The material on the front face of the dike was intended to be sacrificial, and to provide a place for vegetation to establish. I have been monitoring the dike very closely, and vegetation has developed very nicely in most areas. The spot areas where some erosion has occurred are not, at this point, endangering the dike.

In very simple terms, there are two ways an earth fill dike or dam can fail. One is a slope failure and the other is a piping failure. A slope failure occurs when the weight of saturated soil exceeds the internal friction of the soil and a failure plane is created whereby the soil sloughs. This type of failure can be noted to a limited extent on the front face of the dike where erosion has resulted in a near vertical face in a few localized areas, but does not affect the structural portion of the dike. There are no slope failures on the downstream face of the dike, which is to be expected since the dike was built with very stable 6:1 slopes. For a piping failure the phreatic (water) surface must exit on the downstream face of the

dam in order for the dike material to be "piped" through the fill. Since the hydraulic head between the pond and the downstream water surface is only one to two feet, a piping failure is virtually impossible.

The only way that the dike could fail in my opinion would be for erosion to virtually wash away the entire dike. I have many years of experience in the design and analysis of earth fill dams (BLM, Los Angeles County Flood Control District, dam safety inspections for DNRC) and I can assure you that this dike is not in any danger of failing. Since I have stated this opinion many times I can only conclude you do not consider my opinion credible. Therefore I suggest you contact Michelle LeMieaux, Dam Safety Supervisor with the DNRC Dam Safety Bureau for an independent opinion. She is competent in slope stability analysis and dam safety.

You asked that I submit a plan and a schedule for corrective actions. Following is my proposed plan of action:

1) Immediately drop the water level of the pond. By dropping the water level of the pond any erosion of the dike will decline since wave action on the dike will be reduced. The contracted wetland was for 52 acres and the most recent Monitoring Reports prepared by Land and Water Consultants indicated 107 acres of wetland of additional wetland have been created, with additional acreage likely to form with additional time and more normal precipitation. The wetland should be well within the contracted acreage even with this drop in water level. I will remove dam boards and drop the water level on my next trip to the Beaverhead sometime in late January or early February.

2) Provide filter fabric and fill at selected locations. I will inspect and mark the locations where erosion is pronounced, and will cover these areas with filter fabric and gravel. This is cosmetic only, but I realize that you and the Corps will be looking for some action before allowing further credits to be withdrawn. Unless the area is wet and inaccessible I will have this done by May 15, 2004.

I am uncertain as to whether the MDT or the Corps will then inspect the dike and make a judgment as to whether it is to their satisfaction, or if it will then be subject to another personal opinion as to the safety of the dike. If that is the case I would like the Department and/or the Corps to specify what they will accept so I don't have to revisit the site with construction equipment. Contrary to your statement that repair costs would escalate if the structure fails the cost is not in the work, which is minor, but in moving material and equipment to and from the site. I need to know what "look" will appear to be safe to you so I can achieve that appearance.

I am uncomfortable with the awkward relations this has caused between the Department and myself. I again offer to the Department the following options:

a) The Department can take over management of the wetland and manage it any way they see fit. That way the MDT can prohibit grazing, fence in any manner you choose, riprap the dike face, lower or raise the water level to adjust the wetland credits, or whatever

is your pleasure. Since it is apparent we don't agree on management philosophy this is the best option in my opinion.

b) You can allow me to manage the wetland and just provide me with any requirements from the Corps that might jeopardize the credits, which I would take care of immediately. This is how our original agreement was structured and what I understand our agreement to be.

c) I can buy back the easement from the Department and the Department can use the money to fulfill their wetland obligations elsewhere.

Any of these options would be fine by me. The only option that is unacceptable is for the MDT to direct me on how to manage the wetland, but still have me take full responsibility whether or not your directives work. I have been in that position before and it is untenable. If you don't want to manage the wetland yourself, I am agreeable to having a third party manage the wetland according to your directives.

Since it appears unlikely that we will continue with negotiations on the additional wetland credits I would like resolution on all issues at this time. As Mr. Galt is aware, I don't like conflict, and since we will be operating under the original agreement I would like to firmly establish our respective responsibilities so we don't have any more disagreements.

I apologize for the tone of this letter, but you must understand that I have many years of experience in engineering and construction, as well as several other businesses, and have learned to make and trust my own decisions and judgments. I was able to acquire land, negotiate a contract, design the wetland, construct the wetland, and obtain wetland credits in approximately seven months. Now that the project is successful, with credits far in excess of the contract requirements, the Department seems to see a need to micromanage the wetland.


I am also disturbed that when I requested corrections to the Monitoring Report because of factual inaccuracies, you responded that since the accuracies did not affect the conclusions of the report it would not accomplish any purpose to correct them. As you recall, one of the inaccuracies in the report is on page 6 where it states: "Water levels should have been higher, but **the landowner lowered water levels to save the dike** in May 2002." I have no idea where that statement came from, but it is a complete falsehood.

If the Corps is relying on this information to make a judgment on the erosion of the dike, which I would be doing in their position, then I am upset that the record was not set straight. I believe this to be one of the inaccuracies where it does affect the conclusions of the report and the wetland credits. Your refusal to correct this statement may well have been a factor in the determination by the Corps to consider the dike unsafe.

I know this response goes beyond the scope of your letter on the erosion of the dike, but because of the distribution of your copies I feel compelled to state the background and reasoning behind my response. Also I need to know if the actions I propose are acceptable to you and the Corps, and which of the options on management you would like to pursue.

Perhaps an onsite visit with the Department and the Corps would be a good idea so we can be sure that any action taken will be approved. I am looking forward to your response. If you have questions please call me at 447-5000.

Sincerely,



Robert Peccia, P.E.

cc: Allen Steinle – Army Corps  
Dave Galt – Director  
Joel Marshik, P.E. – Chief Engineer  
Dave Hill – Environmental Services  
File



## Dam Safety Program

# Memo

**To:** Bonnie Steg, MDOT, Resources Section Supervisor, Environmental Services,  
**From:** Michele Lemieux, P.E., Dam Safety Program Manager  
**CC:** Timothy McNaboe, Wetland Engineer, Environmental Services  
Bob Peccia, Robert Peccia and Associates  
**Date:** Tuesday, August 02, 2005  
**Re:** Beaverhead Gateway Ranch Wetlands Dike – Trip Report

---

On Tuesday, July 26, 2005 I visited the Beaverhead Gateway Ranch Wetlands Mitigation Project. The purpose of the visit was to evaluate the condition of the upstream face of the dike.

Erosion and beaching due to wave action is present at several locations. In general, the erosion is minor. There are 2 locations where the erosion is beginning to encroach on the crest of the dike: at station 18+00 and at station 13+00 (station locations approximated by pacing).

Erosion of the crest becomes a safety concern when the crest width is narrowed. An embankment with a narrow crest is more susceptible to failure from overtopping. In addition, upstream or downstream slope movement is more apt to cause a failure when the embankment has a narrow crest.

Although the erosion is slowly moving toward the crest of the dike, I consider this to be a maintenance issue and not a safety concern, for several reasons:

First, the eroded areas are still 3 or more feet from the crest of the dike. The crest is very wide and both upstream and downstream slopes are gentle. It would take a considerable amount of additional erosion before the crest would become narrow enough to be a concern.

Second, even with a narrow crest, the dike is not susceptible to failure. The reservoir is located offstream. It is unlikely that an extreme storm event could cause the reservoir level to rise to the point where the dike would be overtopped. In addition, the dike height is low, and the upstream and downstream slopes very gentle. Embankments with this configuration are very stable. Slope failure is unlikely.

Third, the dike was constructed primarily with fat clays. Fat clays are generally non-erosive and resistant to failure.

I would recommend that the dike owner periodically add fill to the areas of the dike where beaching is taking place and vertical slopes are developing. The upstream face should be annually monitored, and a regular maintenance plan developed.



Figure 1. Erosion of upstream face, slowly moving toward crest of dike approximately 375 feet west of overflow structure. Note lack of vegetation on vertical slopes.



Figure 2. View of upstream face looking east. Note gentle slopes.



Figure 3. View of upstream face erosion approximately 870 feet west of overflow structure. Vegetation has a difficult time establishing on vertical slopes, so erosion will likely continue. Crest is still 3 feet away, and very wide in this location, so the erosion is not a threat to the safety and stability of the dam at this time.



Figure 4. Gentle downstream slopes make for a failure resistant structure.

## **Appendix E**

---

### **BIRD SURVEY PROTOCOL GPS PROTOCOL**

---

*MDT Wetland Mitigation Monitoring  
Beaverhead Gateway  
Dillon, Montana*

## **BIRD SURVEY PROTOCOL**

The following is an outline of the MDT Wetland Mitigation Site Monitoring Bird Survey Protocol. Though each site is vastly different, the bird survey data collection methods must be standardized to a certain degree to increase repeatability. An Area Search within a restricted time frame will be used to collect the following data: a bird species list, density, behavior, and habitat-type use. There will be some decisions that team members must make to fit the protocol to their particular site. Each of the following sections and the desired result describes the protocol established to reflect bird species use over time.

### **Species Use within the Mitigation Wetland: Survey Method**

Result: To conduct a bird survey of the wetland mitigation site within a restricted period of time and the budget allotment.

#### ***Sites that can be circumambulated or walked throughout.***

These types of sites will include ponds, enhanced historic river channels, wet meadows, and any area that can be surveyed from the entirety of its perimeter or walked throughout. If the wetland is not uncomfortably inundated, conduct several “meandering” transects through the site in an orderly fashion (record the number and approximate location/direction of the transects in the field notebook; they do not have to be formalized or staked). If a very small portion of the site cannot be crossed due to inundation, this method will also apply. Though the sizes of the site vary, each site will require surveying to the fullest extent possible within a set time limit. The optimum times to conduct the survey are in the morning hours. Conduct the survey from sunrise to no later than 11:00 AM. (Note: some sites may have to be surveyed in the late afternoon or evening due to time constraints or weather; if this is the case, record the time of day and include this information in your report discussion.) If the survey is completed before 11:00 AM and no additions are being made to the list, then the task is complete. The overall limiting factor regarding the number of hours that are spent conducting this survey is the number of budgeted hours; this determination must be made by site by each individual.

In many cases, binoculars will be the only instrument that is needed to identify and count the birds using the wetland. If the wetland includes deep water habitat that can not be assessed with binoculars, then a scope and tripod are necessary. If this is the case, establish as many lookout posts as necessary from key vantage points to collect the data. Depending on the size of the open water, more time may be spent viewing the mitigation area from these vantage points than is spent walking the peripheries of more shallow-water wetlands.

#### ***Sites that cannot be circumambulated.***

These types of sites will include large-bodied waters, such as reservoirs, particularly those with deep water habitat (>6 ft) close to the shore and no wetland development in that area of the shoreline. If one area of the reservoir was graded in such a way to create or enhance the development of a wetland, then that will be the area in which the ambulatory bird survey is conducted. The team member must then determine the length of the shoreline that will be surveyed during each visit.

As stated above in the ambulatory site section, these large sites most likely will have to be surveyed from established vantage points.

### **Species Use within the Mitigation Wetland: Data Recording**

Result: A complete list of bird species using the site, an estimate of bird densities and associated behaviors, and identification of habitat use.

#### ***1. Bird Species List***

Record the bird species on the Bird Survey - Field Data Sheet using the appropriate 4-letter code of the common name. The coding uses the first two letters of the first two words of the birds' common name or if one name, the first four (4) letters. For example, mourning dove is coded MODO and mallard is MALL. If an unknown individual is observed, use the following protocol and define your abbreviation at the bottom of the field data sheet: unknown shorebird: UNSB; unknown brown bird (UNBR); unknown warbler (UNWA); unknown waterfowl (UNWF). For a flyover of a flock of unknown species, use a term that describes the birds' general characteristics and include the approximate flock size in parentheses; do not fill in the habitat column. For example, a flock of black, medium-sized birds could be coded: UNBB / FO (25). You may also note on the data sheet if that particular individual is using a constructed nest box.

#### ***2. Bird Density***

In the office, sum the Bird Survey – Field Data Sheet data by species and by behavior. Record this data in the Bird Summary Table.

#### ***3. Bird Behavior***

Bird behavior must be identified by what is known. When a species is simply observed, the behavior that it is immediately exhibiting is what is recorded. Only behaviors that have discreet descriptive terms should be used. The following terms are recommended: breeding pair individual (BP); foraging (F); flyover (FO); loafing (L; e.g. sleeping, roosting, floating with head tucked under wing are loafing behaviors); and, nesting (N). If more behaviors are observed that do have a specific descriptive word, use them and we will add it to the protocol; descriptive words or phrases such as “migrating” or “living on site” are unknown behaviors.

#### ***4. Bird Species Habitat Use***

We are interested in what bird species are using which particular habitat within the mitigation wetlands. This data is easily collected by simply recording what habitat the species was initially observed. Use the following broad category habitat classifications: aquatic bed (AB - rooted floating, floating-leaved, or submergent vegetation); forested (FO); marsh (MA – cattail, bulrush, emergent vegetation, etc. with surface water); open water (OW – primarily unvegetated); scrub-shrub (SS); and upland buffer (UP); wet meadow (WM – sedges, rushes, grasses with little to no surface water). If other categories are observed onsite that are not suggested here, we will make a new category next year.

## **GPS MAPPING AND AERIAL PHOTO REFERENCING PROCEDURE**

The wetland boundaries, photograph location points and sampling locations were field located with mapping grade Trimble Geo III GPS units. The data was collected with a minimum of three positions per feature using Course/Acquisition code. The collected data was then transferred to a PC and differentially corrected to the nearest operating Community Base Station. The corrected data was then exported to ACAD drawings in Montana State Plain Coordinates NAD 83 international feet.

The GPS positions collected and processed had a 68% accuracy of 7 feet except in isolated areas of Tasks .008 and .011, where it went to 12 feet. This is within the 1 to 5 meter range listed as the expected accuracy of the mapping grade Trimble GPS.

Aerial reference points were used to position the aerial photographs. This positioning did not remove the distortion inherent in all photos; this imagery is to be used as a visual aide only. The located wetland boundaries were given a final review by the wetland biologist and adjustments were made if necessary.

Any relationship of features located to easement or property lines are not to be construed from these figures. These relationships can only be determined with a survey by a licensed surveyor.

## **Appendix F**

---

### **2006 MACROINVERTEBRATE SAMPLING PROTOCOL AND DATA ANALYSIS**

---

*MDT Wetland Mitigation Monitoring  
Beaverhead Gateway  
Dillon, Montana*

# AQUATIC INVERTEBRATE SAMPLING PROTOCOL

## Equipment List

- D-frame sampling net with 1 mm mesh. Wildco is a good source of these.
- Spare net.
- 1-liter plastic sample jars, wide-mouth. VWR has these: catalog #36319-707.
- 95% ethanol: Northwest Scientific in Billings carries this.

All these other things are generally available at hardware or sporting goods stores. Make the labels on an ink jet printer preferably.

- hip waders.
- pre-printed sample labels (printed on Rite-in-the-Rain or other coated paper, two labels per sample).
- pencil.
- plastic pail (3 or 5 gallon).
- large tea strainer or framed screen.
- towel.
- tape for affixing label to jar.
- cooler with ice for sample storage.

## Site Selection

Select the sampling site with these considerations in mind:

- Select a site accessible with hip waders. If substrates are too soft, lay a wide board down to walk on.
- Determine a location that is representative of the overall condition of the wetland.

## Sampling

Wetland invertebrates inhabit the substrate, the water column, the stems and leaves of aquatic vegetation, and the water surface. Your goal is to sweep the collecting net through each of these habitat types, and then to combine the resulting samples into the 1-liter sample jar.

Dip out about a gallon of water into the pail. Pour about a cup of ethanol into the sample jar. Fill out the top half of the sample labels, using pencil, since ink will dissolve in the ethanol.

Ideally, you can sample a swath of water column from near-shore outward to a depth of approximately 3 feet with a long sweep of the net, keeping the net at about half the depth of the water throughout the sweep. Sweep the water surface as well. Pull the net through a vegetated area, beneath the water surface, for at least a meter of distance.

Sample the substrate by pulling the net along the bottom, bumping it against the substrate several times as you pull.

This step is optional, but it gives you a chance to see that you've collected some invertebrates. Rinse the net out into the bucket, and look for insects, crustaceans, etc. If necessary, repeat the sampling process in a nearby location, and add the net contents to the bucket. Remember to sample all four environments.

Sieve the contents of the bucket through the straining device and pour or carefully scrape the contents of the strainer into the sample jar.

If you skip the bucket-and-sieve steps, simply lift handfuls of material out of the sampling net into the jars. In either case, please include some muck or mud and some vegetation in the jar. Often, you will have collected a large amount of vegetable material. If this is the case, lift out handfuls of material from the sieve into the jar, until the jar is about half full. Please limit material you include in the sample, so that there is only a single jar for each sample.

Top off the sample jar with enough ethanol to cover all the material in the jar. Leave as little headroom as possible.

It is not necessary to sample habitats in any specified order. Keep in mind that disturbing the habitats prior to sampling will chase off the animals you are trying to capture.

Complete the sample labels. Place one label inside the sample jar and tape the other label securely to the outside of the jar. Dry the jar before attaching the outer label if necessary. In some situations, it may be necessary to collect more than one sample at a site. If you take multiple samples from the same site, clearly indicate this by using individual sample numbers, along with the total number of samples collected at the site (e.g. Sample #3 of 5 total samples).

Photograph the sampled site.

### **Sample Handling/Shipping**

- In the field, keep collected samples cool by storing them in a cooler. Only a small amount of ice is necessary.
- Inventory all samples, preparing a list of all sites and enumerating all samples, before shipping or delivering to the laboratory.
- Deliver samples to Rhithron.

# **MDT Mitigated Wetland Monitoring Project: Aquatic Invertebrate Monitoring Summary 2001 – 2006**

Prepared for PBS&J, Inc.

Prepared by W.Bollman, Rhithron Associates, Inc.

## **INTRODUCTION**

Among other monitoring activities, aquatic invertebrate assemblages were collected at a number of mitigated wetlands throughout Montana. This report summarizes data generated from six years of collection. Over all years of sampling, a total of 182 invertebrate samples were collected. Table 2 summarizes sites and sampling years.

## **METHODS**

### **Sample processing**

Aquatic invertebrate samples were collected at mitigated wetland sites in the summer months of 2001, 2002, 2003, 2004, 2005 and 2006 by personnel of PBS&J, Inc. Sampling procedures utilized were based on the protocols developed by the Montana Department of Environmental Quality (MT DEQ). Sampling consisted of D-frame net sweeps through emergent vegetation (when present), the water column, and over the water surface, and included disturbing and scraping substrates at each sampled site. These sample components were composited and preserved in ethanol at each wetland site. Samples were delivered to Rhithron Associates, Inc. for processing, taxonomic determinations, and data analysis.

At Rhithron's laboratory, Caton subsamplers and stereomicroscopes with 10X magnification were used to randomly select a minimum of 100 organisms from each sample. In some instances, the entire sample contained fewer than 100 organisms; in these cases, all organisms from the sample were taken. Animals were identified to lowest practical taxonomic levels using relevant published resources. Quality control (QC) procedures were applied to sample sorting, taxonomic determinations and enumeration, and data entry. QC statistics are presented in Table 3. The identified samples have been archived at Rhithron's laboratory.

### **Assessment**

The method employed to assess these wetlands is based on an index incorporating a battery of 12 bioassessment metrics or attributes (Table 1) tested and recommended by Stribling et al. (1995) in a report to the Montana Department of Health and Environmental Science. In that study, it was determined that some of the metrics were of limited use in some geographic regions, and for some wetland types. Despite that finding, all 12 metrics are used in this evaluation of mitigated wetlands, since detailed geographic information and wetland classifications were unavailable.

Scoring criteria for metrics were developed by generally following the tactic used by Stribling et al. Boxplots were generated using a statistical software package (Statistica™), and distributions, median values, ranges, and quartiles for each metric were examined. All sites in all years of sampling were used. Camp Creek, which was sampled in 2002, 2003, 2004, 2005 and 2006, and Kleinschmidt Creek, sampled in 2003, 2004, 2005 and 2006, were assessed using the tested metric battery developed for montane streams of Western Montana (Bollman 1998). Invertebrate assemblages at these sites differed from those of the other sites, and suggested montane or foothill stream conditions rather than wetland conditions. For the wetland sites, "optimal" scores were generally those that fell above the 75<sup>th</sup> percentile (for those metrics that decrease in value in response to stress) or below the 25<sup>th</sup> percentile (for metrics that respond to stress by an increase in value) of all scores. Additional scoring ranges were established by bisecting the range below the 75<sup>th</sup> percentile for decreasing scores (or above the 25<sup>th</sup> percentile for increasing scores) into "sub-optimal" and "poor" assessment categories. A score of 5, 3, or 1 was assigned to optimal, sub-optimal, and poor metric performance, respectively. In this way, metric values were translated into normalized metric scores, and scores for all metrics were summed to produce a total bioassessment score. Total bioassessment scores were classified according to a similar process, using the ranges and distributions of total scores for all sites studied in all years.

The purpose of constructing an index from biological attributes or metrics is to provide a means of integrating information to facilitate the determination of whether management action is needed. The nature of the action needed is not determined solely by the index score, however, but by consideration of an

analysis of the component metrics, the taxonomic composition of the assemblages, and other issues. The diagnostic functions of the metrics and taxonomic data need more study since our understanding of the interrelationships of natural environmental factors and anthropogenic disturbances is tentative. Thus, the further interpretive remarks accompanying the raw taxonomic and metric data in this summary are offered cautiously. Year-to-year comparisons depend on an assumption that specific sites were revisited in each year, and that equivalent sampling methods were utilized at each site revisit.

### **Bioassessment metrics**

An index based on the performance of 12 metrics was constructed, as described above. Table 2 lists those metrics, describes their calculation and the expected response of each to increased degradation or impairment of the wetland.

In addition to the summed scores of each metric and the associated impairment classification described above, each individual metric informs the bioassessment to some degree. The four richness metrics (Total taxa, POET, Chironomidae taxa, and Crustacea taxa + Mollusca taxa) can be interpreted to express habitat complexity as well as water quality. Complex, diverse habitats consist of variable substrates, emergent vegetation, variable water depths and other factors, and are potential features of long-established stable wetlands with minimal human disturbance. In the study conducted by Stribling et al. (1995), all four richness metrics were found to be significantly associated with water quality parameters including conductance, salinity, and total dissolved solids.

Four composition metrics (%Chironomidae, %Orthoclaadiinae of Chironomidae, %Crustacea + %Mollusca, and %Amphipoda) measure the relative contributions of certain taxonomic groups that may have significant responses to habitat and/or water quality impacts. For example, amphipods have been demonstrated to increase in abundance in alkaline conditions. Short-lived, relatively mobile taxa such as chironomids dominate ephemeral environments; many are hemoglobin-bearers capable of tolerating de-oxygenated conditions.

Two tolerance metrics (the Hilsenhoff Biotic Index and %Dominant taxon) were included in the bioassessment battery. The HBI indicates the overall invertebrate assemblage tolerance to nutrient enrichment, warm water, and/or low dissolved oxygen conditions. The percent abundance of the dominant taxon has been demonstrated to be strongly associated with pH, conductance, salinity, total organic carbon, and total dissolved solids.

Two trophic measures (%Collector-gatherers and %Filterers) may be helpful in expressing functional integrity of the invertebrate assemblage, which can be impacted by poor water quality or habitat degradation. High proportions of filtering organisms suggest nutrient and/or organic enrichment, while abundant collectors suggest more positive functional conditions and well-developed wetland morphology. These organisms graze periphyton growing on stable surfaces such as macrophytes.

Metric scoring criteria were re-examined each year as new data was added. For 2005, all 151 records were utilized. Ranges of individual metrics, as well as median metric values remained remarkably consistent over all 5 years of analysis. Since metric value distributions changed insignificantly with the addition of the 2006 data, no changes were made to scoring criteria this year. Summary metric values and scores for the 2006 samples are given in Tables 3a-3d.

### **Quality control**

Quality control procedures for initial sample processing and subsampling involved checking sorting efficiency. These checks were conducted on 100% of the samples by independent technicians who microscopically re-examined 20% of sorted substrate from each sample. All organisms that were missed were counted and this number was added to the total number obtained in the original sort. Sorting efficiency was evaluated by applying the following calculation:

$$SE = \frac{n_1}{n_2} \times 100$$

Where: SE is the sorting efficiency, expressed as a percentage,  $n_1$  is the total number of specimens in the first sort, and  $n_2$  is the total number of specimens in the first and second sorts combined.

Quality control procedures for taxonomic determinations involved checking accuracy, precision and enumeration. Four samples were randomly selected and all organisms re-identified by independent taxonomists. A Bray-Curtis similarity statistic (Bray and Curtis 1957) was generated to evaluate identifications.

**Table 1.** Montana Department of Transportation Mitigated Wetlands Monitoring Project sites, 2001 – 2006.

Site identifier	2001	2002	2003	2004	2005	2006
Beaverhead 1	+	+	+	+	+	+
Beaverhead 2	+	+				
Beaverhead 3	+	+		+	+	+
Beaverhead 4	+	+	+			
Beaverhead 5	+	+	+	+	+	+
Beaverhead 6	+	+	+	+	+	+
Big Sandy 1	+					
Big Sandy 2	+					
Big Sandy 3	+					
Big Sandy 4	+					
Johnson-Valier	+					
VIDA	+					
Cow Coulee	+	+	+			
Fourchette – Puffin	+	+	+	+		
Fourchette – Flashlight	+	+	+	+		
Fourchette – Penguin	+	+	+	+		
Fourchette – Albatross	+	+	+	+		
Big Spring	+	+	+	+	+	
Vince Ames	+					
Ryegate	+					
Lavinia	+					
Stillwater	+	+	+	+	+	
Roundup	+	+	+	+	+	+
Wigeon	+	+	+	+	+	+
Ridgeway	+	+	+	+	+	+
Musgrave – Rest. 1	+	+	+	+	+	+
Musgrave – Rest. 2	+	+	+	+	+	+
Musgrave – Enh. 1	+	+	+	+	+	+
Musgrave – Enh. 2	+					+
Hoskins Landing		+	+	+	+	
Hoskins Landing						
Peterson - 1		+	+	+	+	+
Peterson – 2		+		+	+	+
Peterson – 4		+	+	+	+	+
Peterson – 5		+	+	+	+	+
Jack Johnson - main		+	+			
Jack Johnson - SW		+	+			
Creston		+	+	+	+	
Lawrence Park		+				
Perry Ranch		+			+	
SF Smith River		+	+	+	+	+
Camp Creek		+	+	+	+	+
Camp Creek						+
Kleinschmidt		+	+	+	+	+
Kleinschmidt – stream			+	+	+	+
Ringling - Galt			+			
Circle				+		
Cloud Ranch Pond				+	+	
Cloud Ranch Stream				+		
American Colloid				+	+	+
Jack Creek				+	+	
Jack Creek						
Norem				+	+	+
Rock Creek Ranch					+	+
Wagner Marsh					+	+
Alkali Lake 1						+
Alkali Lake 2						+

**Table 2.** Aquatic invertebrate metrics employed in the MTDT mitigated wetland monitoring study, 2001-2005.

<b>Metric</b>	<b>Metric calculation</b>	<b>Expected response to degradation or impairment</b>
Total taxa	Count of unique taxa identified to lowest recommended taxonomic level	Decrease
POET	Count of unique Plecoptera, Trichoptera, Ephemeroptera, and Odonata taxa identified to lowest recommended taxonomic level	Decrease
Chironomidae taxa	Count of unique midge taxa identified to lowest recommended taxonomic level	Decrease
Crustacea taxa + Mollusca taxa	Count of unique Crustacea taxa and Mollusca taxa identified to lowest recommended taxonomic level	Decrease
% Chironomidae	Percent abundance of midges in the subsample	Increase
Orthoclaadiinae/Chironomidae	Number of individual midges in the sub-family Orthoclaadiinae / total number of midges in the subsample.	Decrease
% Amphipoda	Percent abundance of amphipods in the subsample	Increase
%Crustacea + %Mollusca	Percent abundance of crustaceans in the subsample plus percent abundance of molluscs in the subsample	Increase
HBI	Relative abundance of each taxon multiplied by that taxon's modified Hilsenhoff Biotic Index (tolerance) value. These numbers are summed over all taxa in the subsample.	Increase
%Dominant taxon	Percent abundance of the most abundant taxon in the subsample	Increase
%Collector-Gatherers	Percent abundance of organisms in the collector-gatherer functional group	Decrease
%Filterers	Percent abundance of organisms in the filterer functional group	Increase

## RESULTS

(Note: Individual site discussions were removed from this report by PBS&J and are included in the macroinvertebrate sections of individual monitoring reports. Summary tables (4a – 4d) are provided on the following pages.)

### Quality Assurance

Table 3 gives the results of quality assurance procedures for sample sorting and taxonomic determinations and enumeration.

**Table 3.** Results of quality control procedures for subsampling and taxonomy.

Sample ID	Site name	SE	Bray-Curtis similarity
MDT06PBSJ001	MUSGRAVE LAKE ES-1	91.67%	
MDT06PBSJ002	MUSGRAVE LAKE ES-2	94.44%	
MDT06PBSJ003	MUSGRAVE LAKE RS-1	87.30%	
MDT06PBSJ004	MUSGRAVE LAKE RS-2	100.00%	
MDT06PBSJ005	ROCK CREEK RANCH	96.49%	95.25%
MDT06PBSJ006	Alkali Lake Sample 1	100.00%	
MDT06PBSJ007	Alkali Lake Sample 2	100.00%	
MDT06PBSJ008	Peterson Ranch Pond # 4	100.00%	
MDT06PBSJ009	Peterson Ranch Pond # 1	97.35%	
MDT06PBSJ010	Peterson Ranch Pond # 5	91.67%	
MDT06PBSJ011	South Fork Smith River	100.00%	
MDT06PBSJ012	Beaverhead 1	100.00%	
MDT06PBSJ013	Beaverhead 3	95.65%	
MDT06PBSJ014	Beaverhead 5	100.00%	
MDT06PBSJ015	Beaverhead 6	94.12%	98.38%
MDT06PBSJ016	Peterson Ranch Pond # 2	91.67%	99.66%
MDT06PBSJ017	American Colloid	100.00%	
MDT06PBSJ018	Norem	100.00%	
MDT06PBSJ019	Cloud Ranch	85.56%	98.89%
MDT06PBSJ020	Jack Creek Pond	100.00%	
MDT06PBSJ021	Jack Creek Stream	100.00%	
MDT06PBSJ022	Camp Creek 1	99.10%	
MDT06PBSJ023	Camp Creek 2	100.00%	
MDT06PBSJ024	Kleinschmidt Pond	100.00%	
MDT06PBSJ025	Kleinschmidt Stream	96.49%	
MDT06PBSJ026	Hoskins Landing 1	97.35%	
MDT06PBSJ027	Hoskins Landing 2	96.49%	
MDT06PBSJ028	Wagner Marsh	100.00%	
MDT06PBSJ029	Wigeon Reservoir	100.00%	
MDT06PBSJ030	Ridgeway	98.21%	
MDT06PBSJ031	Roundup	100.00%	

**Table 4a.** Metric values and scores for Montana Department of Transportation mitigated wetland sites. 2006.

	BEAVERHEAD #1	BEAVERHEAD #3	BEAVERHEAD #5	BEAVERHEAD #6	ROUNDUP	WIDGEON	RIDGEWAY	MUSGRAVE RS-1
Total taxa	12	11	4	15	11	11	21	23
POET	1	0	1	3	2	1	3	4
Chironomidae taxa	5	3	1	7	4	3	10	7
Crustacea + Mollusca	1	4	2	3	2	2	5	7
% Chironomidae	52.38%	25.22%	0.69%	63.06%	18.87%	6.42%	37.25%	9.62%
Orthoclaadiinae/Chir	0.181818	0.965517	0	0.142857	0.2	0.285714	0.289474	0.7
% Amphipoda	0.00%	0.00%	0.00%	0.90%	0.00%	6.42%	11.76%	1.92%
% Crustacea + % Mollusca	9.52%	69.57%	98.62%	3.60%	73.58%	79.82%	45.10%	51.92%
HBI	7.857143	7.773913	7.97931	7.243243	8.09434	8.100917	7.127451	7.403846
% Dominant taxon	33.33%	39.13%	97.93%	27.93%	72.64%	73.39%	28.43%	23.08%
% Collector-Gatherers	61.90%	68.70%	100.00%	84.68%	87.74%	6.42%	49.02%	47.12%
% Filterers	0.00%	2.61%	0.00%	1.80%	0.00%	0.00%	0.00%	4.81%
Total taxa	1	1	1	3	1	1	5	5
POET	1	1	1	3	1	1	3	5
Chironomidae taxa	3	3	1	5	3	3	5	5
Crustacea + Mollusca	1	3	1	1	1	1	3	5
% Chironomidae	1	3	5	1	3	5	3	5
Orthoclaadiinae/Chir	1	5	1	1	3	3	3	5
% Amphipoda	5	5	5	5	5	3	3	5
% Crustacea + % Mollusca	5	1	1	5	1	1	3	3
HBI	1	1	1	3	1	1	3	3
% Dominant taxon	5	3	1	5	1	1	5	5
% Collector-Gatherers	3	3	5	5	5	1	3	3
% Filterers	3	3	3	3	3	3	3	3
<b>Total score</b>	30	32	26	40	28	24	42	52
<b>Percent of maximum score</b>	0.5	0.533333	0.433333	0.666667	0.466667	0.4	0.7	0.866667
<b>Impairment classification</b>	poor	poor	poor	sub-optimal	poor	poor	optimal	optimal

**Table 4b.** Metric values and scores for Montana Department of Transportation mitigated wetland sites. 2006.

	MUSGRAVE RS- 2	MUSGRAVE ES- 1	MUSGRAVE ES- 2	HOSKINS LANDING 1	HOSKINS LANDING 2	PETERSON RANCH 1	PETERSON RANCH 2	PETERSON RANCH 4	PETERSON RANCH 5
<b>Total taxa</b>	10	21	10	22	29	19	17	28	26
POET	1	2	1	5	4	2	2	3	4
Chironomidae taxa	2	7	4	6	6	7	4	13	9
Crustacea + Mollusca	3	6	0	5	9	5	6	5	6
% Chironomidae	3.96%	10.89%	10.00%	18.18%	11.71%	64.08%	7.48%	27.52%	14.29%
Orthoclaadiinae/Chir	0	0.181818	0.125	0.055556	0.307692	0.757576	0.75	0.6	0.75
% Amphipoda	0.00%	2.97%	0.00%	5.05%	1.80%	1.94%	22.43%	2.75%	15.18%
% Crustacea + % Mollusca	8.91%	75.25%	0.00%	20.20%	23.42%	8.74%	42.06%	19.27%	40.18%
HBI	6.326733	6.940594	6	7.111111	7.585586	6.631068	6.719626	7.293578	7.321429
% Dominant taxon	70.30%	38.61%	83.75%	25.25%	42.34%	47.57%	28.04%	20.18%	16.07%
% Collector-Gatherers	15.84%	8.91%	3.75%	64.65%	62.16%	72.82%	31.78%	34.86%	50.89%
% Filterers	0.00%	0.00%	0.00%	6.06%	5.41%	3.88%	3.74%	8.26%	0.89%
<b>Total taxa</b>	1	5	1	5	5	3	3	5	5
POET	1	1	1	5	5	1	1	3	5
Chironomidae taxa	1	5	3	3	3	5	3	5	5
Crustacea + Mollusca	1	5	1	3	5	3	5	3	5
% Chironomidae	5	5	5	3	5	1	5	3	5
Orthoclaadiinae/Chir	1	1	1	1	3	5	5	5	5
% Amphipoda	5	5	5	3	5	5	3	5	3
% Crustacea + % Mollusca	5	1	5	5	5	5	3	5	3
HBI	5	3	5	3	3	5	5	3	3
% Dominant taxon	1	3	1	5	3	3	5	5	5
% Collector-Gatherers	1	1	1	3	3	3	1	1	3
% Filterers	3	3	3	1	3	3	3	1	3
<b>Total score</b>	30	38	32	40	48	42	42	44	50
<b>Percent of maximum score</b>	0.5	0.633333	0.533333	0.666667	0.8	0.7	0.7	0.733333	0.833333
<b>Impairment classification</b>	<b>poor</b>	<b>sub-optimal</b>	<b>poor</b>	<b>sub-optimal</b>	<b>optimal</b>	<b>optimal</b>	<b>optimal</b>	<b>optimal</b>	<b>optimal</b>

**Table 4c.** Metric values and scores for Montana Department of Transportation mitigated wetland sites. 2006

	<b>SOUTH FORK SMITH RIVER</b>	<b>CAMP CREEK 1*</b>	<b>CAMP CREEK 2*</b>	<b>KLEINSCH MIDT POND</b>	<b>KLEINSCH MIDT STREAM*</b>	<b>CLOUD RANCH</b>	<b>COLLOID</b>	<b>JACK CREEK POND</b>	<b>JACK CREEK STREAM</b>
Total taxa	14	31	29	20	22	13	7	7	5
POET	4	8	8	5	1	1	2	0	0
Chironomidae taxa	3	10	8	6	8	6	4	4	0
Crustacea + Mollusca	4	1	3	2	5	3	0	2	2
% Chironomidae	18.02%	45.87%	16.07%	8.04%	77.68%	23.81%	84.21%	75.00%	0.00%
Orthoclaadiinae/Chir	0.05	0.26	0.277778	0.222222	0.448276	0.65	0.25	0.555556	0
% Amphipoda	18.02%	0.00%	0.00%	25.00%	0.00%	4.76%	0.00%	0.00%	5.00%
% Crustacea + % Mollusca	58.56%	0.92%	3.57%	25.89%	5.36%	11.90%	0.00%	16.67%	7.50%
HBI	7.540541	4.504587	4.294643	7.241071	5.928571	7.535714	6.315789	8.833333	7.325
% Dominant taxon	25.23%	24.77%	37.50%	25.00%	33.93%	36.90%	52.63%	33.33%	60.00%
% Collector-Gatherers	41.44%	48.62%	31.25%	62.50%	46.43%	64.29%	21.05%	58.33%	67.50%
% Filterers	15.32%	6.42%	7.14%	3.57%	38.39%	2.38%	0.00%	0.00%	0.00%
Total taxa	1	5	5	3	5	1	1	1	1
POET	5	5	5	5	1	1	1	1	1
Chironomidae taxa	3	5	5	3	5	3	3	3	1
Crustacea + Mollusca	3	1	1	1	3	1	1	1	1
% Chironomidae	3	1	5	5	1	3	1	1	5
Orthoclaadiinae/Chir	1	3	3	3	3	5	3	5	1
% Amphipoda	3	5	5	1	5	3	5	5	3
% Crustacea + % Mollusca	3	5	5	5	5	5	5	5	5
HBI	3	5	5	3	5	3	5	1	3
% Dominant taxon	5	5	3	5	5	3	1	5	1
% Collector-Gatherers	1	3	1	3	3	3	1	3	3
% Filterers	1	1	1	3	1	3	3	3	3
<b>Total score</b>	32	44	44	40	42	34	30	34	28
<b>Percent of maximum score</b>	0.533333	0.733333	0.733333	0.666667	0.7	0.566667	0.5	0.566667	0.466667
<b>Impairment classification</b>	<b>poor</b>	<i>optimal</i>	<i>optimal</i>	<i>sub-optimal</i>	<i>optimal</i>	<i>sub-optimal</i>	<b>poor</b>	<i>sub-optimal</i>	<b>poor</b>

\*Sites indicated by asterisks were dominated by lotic fauna, and were evaluated with the MDEQ index for streams in the text and charts. Scores and impairment classifications in this table (italicized) are included only for completeness and are not reliable indications of conditions at these sites. See text.

**Table 4d.** Metric values and scores for Montana Department of Transportation mitigated wetland sites. 2006.

	NOREM	ROCK CREEK RANCH	WAGNER MARSH	ALKALI LAKE 1	ALKALI LAKE 2
Total taxa	6	15	11	6	5
POET	1	0	0	0	0
Chironomidae taxa	2	4	4	3	0
Crustacea + Mollusca	1	4	3	1	1
% Chironomidae	82.93%	8.40%	13.51%	42.86%	0.00%
Orthoclaadiinae/Chir	0	0.2	0.6	0.666667	0
% Amphipoda	0.00%	0.00%	0.00%	0.00%	0.00%
% Crustacea + % Mollusca	7.32%	65.55%	23.42%	7.14%	9.52%
HBI	7.317073	7.638655	7.036036	7.785714	7.904762
% Dominant taxon	65.85%	47.06%	45.95%	42.86%	52.38%
% Collector-Gatherers	68.29%	56.30%	47.75%	28.57%	9.52%
% Filterers	17.07%	0.00%	0.90%	0.00%	0.00%
Total taxa	1	3	1	1	1
POET	1	1	1	1	1
Chironomidae taxa	1	3	3	3	1
Crustacea + Mollusca	1	3	1	1	1
% Chironomidae	1	5	5	1	5
Orthoclaadiinae/Chir	1	3	5	5	1
% Amphipoda	5	5	5	5	5
% Crustacea + % Mollusca	5	1	5	5	5
HBI	3	1	3	1	1
% Dominant taxon	1	3	3	3	1
% Collector-Gatherers	3	3	3	1	1
% Filterers	1	3	3	3	3
<b>Total score</b>	24	34	38	30	26
<b>Percent of maximum score</b>	0.4	0.566667	0.633333	0.5	0.433333
<b>Impairment classification</b>	<b>poor</b>	<b>sub-optimal</b>	<b>sub-optimal</b>	<b>poor</b>	<b>poor</b>

### **Literature cited**

Bollman, W. 1998. Montana Valleys and Foothill Prairies Ecoregion. Master's Thesis. (M.S.) University of Montana. Missoula, Montana.

Bukantis, R. 1998. Rapid bioassessment macroinvertebrate protocols: Sampling and sample analysis SOP's. Working draft. Montana Department of Environmental Quality. Planning Prevention and Assistance Division. Helena, Montana.

McCune, B. and J.B. Grace. 2002. Analysis of Ecological Communities. MjM Software Design, Gleneden Beach, Oregon, USA.

McCune, B. and M.J. Mefford. 2002. PC-ORD. Multivariate Analysis of Ecological Data, Version 4. MjM Software Design, Gleneden Beach, Oregon, USA.

Stribling, J.B., J. Lathrop-Davis, M.T. Barbour, J.S. White, and E.W. Leppo. 1995. Evaluation of environmental indicators for the wetlands of Montana: the multimetric approach using benthic macroinvertebrates. Report to the Montana Department of Health and Environmental Science. Helena, Montana.

# Taxa Listing

Project ID: MDT06PBSJ  
RAI No.: MDT06PBSJ012

RAI No.: MDT06PBSJ012

Sta. Name: Beaverhead 1

Client ID:

Date Coll.:

No. Jars: 1

STORET ID:

Taxonomic Name	Count	PRA	Unique	Stage	Qualifier	BI	Function
<b>Non-Insect</b>							
Acari	1	4.76%	Yes	Unknown		5	PR
Nematoda	1	4.76%	Yes	Unknown		5	PA
Ostracoda	2	9.52%	Yes	Unknown		8	CG
<b>Ephemeroptera</b>							
Caenidae							
<i>Caenis</i> sp.	1	4.76%	Yes	Larva		7	CG
<b>Heteroptera</b>							
Corixidae							
<i>Palmacorixa</i> sp.	1	4.76%	Yes	Adult		5	PR
<i>Sigara</i> sp.	1	4.76%	Yes	Adult		5	PH
<b>Diptera</b>							
Ceratopogonidae							
Ceratopogoninae	3	14.29%	Yes	Larva		6	PR
<b>Chironomidae</b>							
Chironomidae							
<i>Acricotopus</i> sp.	1	4.76%	Yes	Larva		10	CG
<i>Chironomus</i> sp.	7	33.33%	Yes	Larva		10	CG
Orthoclaadiinae	1	4.76%	No	Larva	Early Instar	6	CG
<i>Paratanytarsus</i> sp.	1	4.76%	Yes	Larva		6	CG
Tanypodinae	1	4.76%	Yes	Larva	Early Instar	7	PR
	<b>Sample Count</b>	<b>21</b>					

# Metrics Report

Project ID: MDT06PBSJ  
 RAI No.: MDT06PBSJ012  
 Sta. Name: Beaverhead 1  
 Client ID:  
 STORET ID:  
 Coll. Date:


## Abundance Measures

Sample Count: 21  
 Sample Abundance: 21.00 100.00% of sample used

Coll. Procedure:  
 Sample Notes:

## Taxonomic Composition

Category	R	A	PRA
Non-Insect	3	4	19.05%
Odonata			
Ephemeroptera	1	1	4.76%
Plecoptera			
Heteroptera	2	2	9.52%
Megaloptera			
Trichoptera			
Lepidoptera			
Coleoptera			
Diptera	1	3	14.29%
Chironomidae	4	11	52.38%




## Dominant Taxa

Category	A	PRA
Chironomus	7	33.33%
Ceratopogoninae	3	14.29%
Ostracoda	2	9.52%
Tanypodinae	1	4.76%
Sigara	1	4.76%
Paratanytarsus	1	4.76%
Palmarcorixa	1	4.76%
Orthoclaadiinae	1	4.76%
Nematoda	1	4.76%
Caenis	1	4.76%
Acricotopus	1	4.76%
Acari	1	4.76%

## Functional Composition

Category	R	A	PRA
Predator	4	6	28.57%
Parasite	1	1	4.76%
Collector Gatherer	5	13	61.90%
Collector Filterer			
Macrophyte Herbivore			
Piercer Herbivore	1	1	4.76%
Xylophage			
Scraper			
Shredder			
Omnivore			
Unknown			




## Metric Values and Scores

Metric	Value	BIBI	MTP	MTV	MTM
<i>Composition</i>					
Taxa Richness	11	1	0		0
Non-Insect Percent	19.05%				
E Richness	1	1		0	
P Richness	0	1		0	
T Richness	0	1		0	
EPT Richness	1		0		0
EPT Percent	4.76%		0		0
Oligochaeta+Hirudinea Percent					
Baetidae/Ephemeroptera	0.000				
Hydropsychidae/Trichoptera	0.000				
<i>Dominance</i>					
Dominant Taxon Percent	33.33%		2		2
Dominant Taxa (2) Percent	47.62%				
Dominant Taxa (3) Percent	57.14%	3			
Dominant Taxa (10) Percent	90.48%				
<i>Diversity</i>					
Shannon H (loge)	2.081				
Shannon H (log2)	3.002		3		
Margalef D	3.338				
Simpson D	0.132				
Evenness	0.106				
<i>Function</i>					
Predator Richness	4		2		
Predator Percent	28.57%	5			
Filterer Richness	0				
Filterer Percent	0.00%			3	
Collector Percent	61.90%		2		2
Scraper+Shredder Percent	0.00%		0		0
Scraper/Filterer	0.000				
Scraper/Scraper+Filterer	0.000				
<i>Habit</i>					
Burrower Richness	2				
Burrower Percent	47.62%				
Swimmer Richness	2				
Swimmer Percent	9.52%				
Clinger Richness	0	1			
Clinger Percent	0.00%				
<i>Characteristics</i>					
Cold Stenotherm Richness	0				
Cold Stenotherm Percent	0.00%				
Hemoglobin Bearer Richness	1				
Hemoglobin Bearer Percent	33.33%				
Air Breather Richness	0				
Air Breather Percent	0.00%				
<i>Voltinism</i>					
Univoltine Richness	4				
Semivoltine Richness	0	1			
Multivoltine Percent	71.43%		1		
<i>Tolerance</i>					
Sediment Tolerant Richness	0				
Sediment Tolerant Percent	0.00%				
Sediment Sensitive Richness	0				
Sediment Sensitive Percent	0.00%				
Metals Tolerance Index	3.867				
Pollution Sensitive Richness	0	1		0	
Pollution Tolerant Percent	38.10%	3		0	
Hilsenhoff Biotic Index	7.619		0		0
Intolerant Percent	0.00%				
Supertolerant Percent	47.62%				
CTQa	103.500				

## Bioassessment Indices

BioIndex	Description	Score	Pct	Rating
BIBI	B-IBI (Karr et al.)	18	36.00%	
MTP	Montana DEQ Plains (Bukantis 1998)	10	33.33%	Moderate
MTV	Montana Revised Valleys/Foothills (Bollman 1998)	3	16.67%	Severe
MTM	Montana DEQ Mountains (Bukantis 1998)	4	19.05%	Severe



# Taxa Listing

Project ID: MDT06PBSJ  
RAI No.: MDT06PBSJ013

RAI No.: MDT06PBSJ013

Sta. Name: Beaverhead 3

Client ID:

Date Coll.:

No. Jars: 1

STORET ID:

Taxonomic Name	Count	PRA	Unique	Stage	Qualifier	BI	Function
<b>Non-Insect</b>							
Cladocera	3	2.61%	Yes	Unknown		8	CF
Copepoda	29	25.22%	Yes	Unknown		8	CG
Nematoda	1	0.87%	Yes	Unknown		5	PA
Ostracoda	45	39.13%	Yes	Unknown		8	CG
Physidae							
Physidae	3	2.61%	Yes	Unknown		8	SC
Tubificidae							
Tubificidae	3	2.61%	Yes	Unknown		10	CG
<b>Coleoptera</b>							
Dytiscidae							
Dytiscidae	1	0.87%	Yes	Larva		5	PR
<b>Diptera</b>							
Syrphidae							
Syrphidae	1	0.87%	Yes	Larva		10	CG
<b>Chironomidae</b>							
Chironomidae							
<i>Cricotopus (Isocladius) sp.</i>	26	22.61%	Yes	Larva		7	SH
<i>Dicrotendipes sp.</i>	1	0.87%	Yes	Larva		8	CG
<i>Phaenopsectra sp.</i>	2	1.74%	Yes	Larva		7	SC
<b>Sample Count</b>	<b>115</b>						

# Metrics Report

Project ID: MDT06PBSJ  
 RAI No.: MDT06PBSJ013  
 Sta. Name: Beaverhead 3  
 Client ID:  
 STORET ID:  
 Coll. Date:


## Abundance Measures

Sample Count: 115  
 Sample Abundance: 1,725.00 6.67% of sample used

Coll. Procedure:  
 Sample Notes:

## Taxonomic Composition

Category	R	A	PRA
Non-Insect	6	84	73.04%
Odonata			
Ephemeroptera			
Plecoptera			
Heteroptera			
Megaloptera			
Trichoptera			
Lepidoptera			
Coleoptera	1	1	0.87%
Diptera	1	1	0.87%
Chironomidae	3	29	25.22%




## Dominant Taxa

Category	A	PRA
Ostracoda	45	39.13%
Copepoda	29	25.22%
Cricotopus (Isocladius)	26	22.61%
Tubificidae	3	2.61%
Physidae	3	2.61%
Cladocera	3	2.61%
Phaenopsectra	2	1.74%
Syrphidae	1	0.87%
Nematoda	1	0.87%
Dytiscidae	1	0.87%
Dicrotendipes	1	0.87%

## Functional Composition

Category	R	A	PRA
Predator	1	1	0.87%
Parasite	1	1	0.87%
Collector Gatherer	5	79	68.70%
Collector Filterer	1	3	2.61%
Macrophyte Herbivore			
Piercer Herbivore			
Xylophage			
Scraper	2	5	4.35%
Shredder	1	26	22.61%
Omnivore			
Unknown			




## Metric Values and Scores

Metric	Value	BIBI	MTP	MTV	MTM
<i>Composition</i>					
Taxa Richness	11	1	0		0
Non-Insect Percent	73.04%				
E Richness	0	1		0	
P Richness	0	1		0	
T Richness	0	1		0	
EPT Richness	0		0		0
EPT Percent	0.00%		0		0
Oligochaeta+Hirudinea Percent	2.61%				
Baetidae/Ephemeroptera	0.000				
Hydropsychidae/Trichoptera	0.000				
<i>Dominance</i>					
Dominant Taxon Percent	39.13%		2		1
Dominant Taxa (2) Percent	64.35%				
Dominant Taxa (3) Percent	86.96%	1			
Dominant Taxa (10) Percent	99.13%				
<i>Diversity</i>					
Shannon H (loge)	1.572				
Shannon H (log2)	2.267		1		
Margalef D	2.108				
Simpson D	0.264				
Evenness	0.132				
<i>Function</i>					
Predator Richness	1		0		
Predator Percent	0.87%	1			
Filterer Richness	1				
Filterer Percent	2.61%			3	
Collector Percent	71.30%		2		1
Scraper+Shredder Percent	26.96%		2		1
Scraper/Filterer	1.667				
Scraper/Scraper+Filterer	0.625				
<i>Habit</i>					
Burrower Richness	1				
Burrower Percent	0.87%				
Swimmer Richness	0				
Swimmer Percent	0.00%				
Clinger Richness	2	1			
Clinger Percent	24.35%				
<i>Characteristics</i>					
Cold Stenotherm Richness	0				
Cold Stenotherm Percent	0.00%				
Hemoglobin Bearer Richness	3				
Hemoglobin Bearer Percent	5.22%				
Air Breather Richness	2				
Air Breather Percent	1.74%				
<i>Voltinism</i>					
Univoltine Richness	3				
Semivoltine Richness	1	1			
Multivoltine Percent	93.04%		0		
<i>Tolerance</i>					
Sediment Tolerant Richness	1				
Sediment Tolerant Percent	2.61%				
Sediment Sensitive Richness	0				
Sediment Sensitive Percent	0.00%				
Metals Tolerance Index	4.727				
Pollution Sensitive Richness	0				
Pollution Tolerant Percent	7.83%	1		0	
Hilsenhoff Biotic Index	7.774		0		0
Intolerant Percent	0.00%				
Supertolerant Percent	73.91%				
CTQa	102.000				

## Bioassessment Indices

BioIndex	Description	Score	Pct	Rating
BIBI	B-IBI (Karr et al.)	14	28.00%	
MTP	Montana DEQ Plains (Bukantis 1998)	7	23.33%	Moderate
MTV	Montana Revised Valleys/Foothills (Bollman 1998)	5	27.78%	Moderate
MTM	Montana DEQ Mountains (Bukantis 1998)	3	14.29%	Severe



# Taxa Listing

Project ID: MDT06PBSJ  
RAI No.: MDT06PBSJ014

RAI No.: MDT06PBSJ014

Sta. Name: Beaverhead 5

Client ID:

Date Coll.:

No. Jars: 1

STORET ID:

Taxonomic Name	Count	PRA	Unique	Stage	Qualifier	BI	Function
<b>Non-Insect</b>							
Copepoda	1	0.69%	Yes	Unknown		8	CG
Ostracoda	142	97.93%	Yes	Unknown		8	CG
<b>Ephemeroptera</b>							
Caenidae							
<i>Caenis</i> sp.	1	0.69%	Yes	Larva		7	CG
<b>Chironomidae</b>							
Chironomidae							
Chironomini	1	0.69%	Yes	Larva	Damaged	6	CG
Sample Count	145						

# Metrics Report

Project ID: MDT06PBSJ  
 RAI No.: MDT06PBSJ014  
 Sta. Name: Beaverhead 5  
 Client ID:  
 STORET ID:  
 Coll. Date:


## Abundance Measures

Sample Count: 145  
 Sample Abundance: 621.43 23.33% of sample used

Coll. Procedure:  
 Sample Notes:

## Taxonomic Composition

Category	R	A	PRA
Non-Insect	2	143	98.62%
Odonata			
Ephemeroptera	1	1	0.69%
Plecoptera			
Heteroptera			
Megaloptera			
Trichoptera			
Lepidoptera			
Coleoptera			
Diptera			
Chironomidae	1	1	0.69%




## Dominant Taxa

Category	A	PRA
Ostracoda	142	97.93%
Copepoda	1	0.69%
Chironomina	1	0.69%
Caenis	1	0.69%

## Functional Composition

Category	R	A	PRA
Predator			
Parasite			
Collector Gatherer	4	145	100.00%
Collector Filterer			
Macrophyte Herbivore			
Piercer Herbivore			
Xylophage			
Scraper			
Shredder			
Omnivore			
Unknown			




## Metric Values and Scores

Metric	Value	BIBI	MTP	MTV	MTM
<i>Composition</i>					
Taxa Richness	4	1	0		0
Non-Insect Percent	98.62%				
E Richness	1	1		0	
P Richness	0	1		0	
T Richness	0	1		0	
EPT Richness	1		0		0
EPT Percent	0.69%		0		0
Oligochaeta+Hirudinea Percent					
Baetidae/Ephemeroptera	0.000				
Hydropsychidae/Trichoptera	0.000				
<i>Dominance</i>					
Dominant Taxon Percent	97.93%		0		0
Dominant Taxa (2) Percent	98.62%				
Dominant Taxa (3) Percent	99.31%	1			
Dominant Taxa (10) Percent	100.00%				
<i>Diversity</i>					
Shannon H (loge)	0.123				
Shannon H (log2)	0.178		0		
Margalef D	0.603				
Simpson D	0.959				
Evenness	0.015				
<i>Function</i>					
Predator Richness	0		0		
Predator Percent	0.00%	1			
Filterer Richness	0				
Filterer Percent	0.00%			3	
Collector Percent	100.00%		0		0
Scraper+Shredder Percent	0.00%		0		0
Scraper/Filterer	0.000				
Scraper/Scraper+Filterer	0.000				
<i>Habit</i>					
Burrower Richness	1				
Burrower Percent	0.69%				
Swimmer Richness	0				
Swimmer Percent	0.00%				
Clinger Richness	0	1			
Clinger Percent	0.00%				
<i>Characteristics</i>					
Cold Stenotherm Richness	0				
Cold Stenotherm Percent	0.00%				
Hemoglobin Bearer Richness					
Hemoglobin Bearer Percent					
Air Breather Richness	0				
Air Breather Percent	0.00%				
<i>Voltinism</i>					
Univoltine Richness	1				
Semivoltine Richness	0	1			
Multivoltine Percent	99.31%		0		
<i>Tolerance</i>					
Sediment Tolerant Richness	0				
Sediment Tolerant Percent	0.00%				
Sediment Sensitive Richness	0				
Sediment Sensitive Percent	0.00%				
Metals Tolerance Index	3.000				
Pollution Sensitive Richness	0	1		0	
Pollution Tolerant Percent	0.69%	5		3	
Hilsenhoff Biotic Index	7.979		0		0
Intolerant Percent	0.00%				
Supertolerant Percent	98.62%				
CTQa	90.000				

## Bioassessment Indices

BiIndex	Description	Score	Pct	Rating
BIBI	B-IBI (Karr et al.)	14	28.00%	
MTP	Montana DEQ Plains (Bukantis 1998)	0	0.00%	Severe
MTV	Montana Revised Valleys/Foothills (Bollman 1998)	6	33.33%	Moderate
MTM	Montana DEQ Mountains (Bukantis 1998)	0	0.00%	Severe



# Taxa Listing

Project ID: MDT06PBSJ  
RAI No.: MDT06PBSJ015

RAI No.: MDT06PBSJ015

Sta. Name: Beaverhead 6

Client ID:

Date Coll.:

No. Jars: 1

STORET ID:

Taxonomic Name	Count	PRA	Unique	Stage	Qualifier	BI	Function
<b>Non-Insect</b>							
Acari	2	1.80%	Yes	Unknown		5	PR
Ostracoda	2	1.80%	Yes	Unknown		8	CG
Turbellaria	2	1.80%	Yes	Unknown		4	PR
Physidae							
Physidae	1	0.90%	Yes	Unknown		8	SC
Talitridae							
<i>Hyalella</i> sp.	1	0.90%	Yes	Unknown		8	CG
<b>Odonata</b>							
Coenagrionidae							
<i>Enallagma</i> sp.	1	0.90%	Yes	Larva		7	PR
<b>Ephemeroptera</b>							
Baetidae							
<i>Callibaetis</i> sp.	1	0.90%	Yes	Larva		9	CG
Caenidae							
<i>Caenis</i> sp.	31	27.93%	Yes	Larva		7	CG
<b>Chironomidae</b>							
Chironomidae							
<i>Apedilum</i> sp.	7	6.31%	Yes	Larva		11	CG
Chironomidae	2	1.80%	No	Pupa		10	CG
<i>Cricotopus (Isocladius)</i> sp.	9	8.11%	Yes	Larva		7	SH
<i>Dicrotendipes</i> sp.	28	25.23%	Yes	Larva		8	CG
Orthoclaadiinae	1	0.90%	No	Larva	Early Instar	6	CG
<i>Paratanytarsus</i> sp.	21	18.92%	Yes	Larva		6	CG
Tanytarsini	2	1.80%	No	Larva	Early Instar	6	CF
<b>Sample Count</b>	<b>111</b>						

# Metrics Report

Project ID: MDT06PBSJ  
 RAI No.: MDT06PBSJ015  
 Sta. Name: Beaverhead 6  
 Client ID:  
 STORET ID:  
 Coll. Date:


## Abundance Measures

Sample Count: 111  
 Sample Abundance: 666.00 16.67% of sample used

Coll. Procedure:  
 Sample Notes:

## Taxonomic Composition

Category	R	A	PRA
Non-Insect	5	8	7.21%
Odonata	1	1	0.90%
Ephemeroptera	2	32	28.83%
Plecoptera			
Heteroptera			
Megaloptera			
Trichoptera			
Lepidoptera			
Coleoptera			
Diptera			
Chironomidae	4	70	63.06%




## Dominant Taxa

Category	A	PRA
Caenis	31	27.93%
Dicrotendipes	28	25.23%
Paratanytarsus	21	18.92%
Cricotopus (Isociadius)	9	8.11%
Apedilum	7	6.31%
Turbellaria	2	1.80%
Tanytarsini	2	1.80%
Ostracoda	2	1.80%
Chironomidae	2	1.80%
Acari	2	1.80%
Physidae	1	0.90%
Orthoclaadiinae	1	0.90%
Hyalella	1	0.90%
Enallagma	1	0.90%
Callibaetis	1	0.90%

## Functional Composition

Category	R	A	PRA
Predator	3	5	4.50%
Parasite			
Collector Gatherer	7	94	84.68%
Collector Filterer	0	2	1.80%
Macrophyte Herbivore			
Piercer Herbivore			
Xylophage			
Scraper	1	1	0.90%
Shredder	1	9	8.11%
Omnivore			
Unknown			



## Metric Values and Scores

Metric	Value	BIBI	MTP	MTV	MTM
<i>Composition</i>					
Taxa Richness	12	1	1		0
Non-Insect Percent	7.21%				
E Richness	2	1		1	
P Richness	0	1		0	
T Richness	0	1		0	
EPT Richness	2		0		0
EPT Percent	28.83%		1		0
Oligochaeta+Hirudinea Percent					
Baetidae/Ephemeroptera	0.031				
Hydropsychidae/Trichoptera	0.000				
<i>Dominance</i>					
Dominant Taxon Percent	27.93%		3		2
Dominant Taxa (2) Percent	53.15%				
Dominant Taxa (3) Percent	72.07%	3			
Dominant Taxa (10) Percent	95.50%				
<i>Diversity</i>					
Shannon H (loge)	1.821				
Shannon H (log2)	2.628		2		
Margalef D	2.359				
Simpson D	0.200				
Evenness	0.120				
<i>Function</i>					
Predator Richness	3		1		
Predator Percent	4.50%	1			
Filterer Richness	0				
Filterer Percent	1.80%			3	
Collector Percent	86.49%		1		0
Scraper+Shredder Percent	9.01%		1		0
Scraper/Filterer	0.500				
Scraper/Scraper+Filterer	0.333				
<i>Habit</i>					
Burrower Richness	1				
Burrower Percent	25.23%				
Swimmer Richness	1				
Swimmer Percent	0.90%				
Clinger Richness	1	1			
Clinger Percent	8.11%				
<i>Characteristics</i>					
Cold Stenotherm Richness	0				
Cold Stenotherm Percent	0.00%				
Hemoglobin Bearer Richness	2				
Hemoglobin Bearer Percent	31.53%				
Air Breather Richness	0				
Air Breather Percent	0.00%				
<i>Voltinism</i>					
Univoltine Richness	4				
Semivoltine Richness	0	1			
Multivoltine Percent	69.37%		1		
<i>Tolerance</i>					
Sediment Tolerant Richness	0				
Sediment Tolerant Percent	0.00%				
Sediment Sensitive Richness	0				
Sediment Sensitive Percent	0.00%				
Metals Tolerance Index	3.443				
Pollution Sensitive Richness	0	1			0
Pollution Tolerant Percent	54.96%	1			0
Hilsenhoff Biotic Index	7.058		0		0
Intolerant Percent	0.00%				
Supertolerant Percent	31.53%				
CTQa	99.000				

## Bioassessment Indices

BiIndex	Description	Score	Pct	Rating
BIBI	B-IBI (Karr et al.)	12	24.00%	
MTP	Montana DEQ Plains (Bukantis 1998)	11	36.67%	Moderate
MTV	Montana Revised Valleys/Foothills (Bollman 1998)	4	22.22%	Moderate
MTM	Montana DEQ Mountains (Bukantis 1998)	2	9.52%	Severe

